

THE ERA KEY  
TO THE  
UNITED STATES  
PHARMACOPŒIA  
1894.

NATIONAL LIBRARY OF MEDICINE  
Bethesda, Maryland

*Gift of*  
William H. Helfand



COMPLIMENTS  
OF  
MELLIN'S FOOD  
BOSTON, MASS.



# MELLIN'S FOOD

—FOR—

INFANTS AND INVALIDS.

---

**Mellin's Food** is entirely free from husks and indigestible inert matter.

**Mellin's Food** is entirely free from animal germs.

**Mellin's Food** is entirely free from cane sugar.

**Mellin's Food** keeps for any length of time in all climates without deterioration or change.

**Mellin's Food** is not a mechanical mixture of cereals.

---

DOLIBER-GOODALE CO.,

BOSTON, MASS.

# MELLIN'S FOOD

— FOR —

INFANTS AND INVALIDS.

---

The mixture of **Mellin's Food** and fresh cow's milk shows the closest approximation, chemically and physiologically, to mother's milk. The **MELLIN'S FOOD** causes the albuminates of the cow's milk to become light and flocculent in the stomach as in mother's milk; otherwise they would be coagulated into a tough, hard curd in the stomach.

**Mellin's Food** is the only infant food which fulfils the requirements of Liebig's principles.

**Mellin's Food** prepared with fresh cow's milk is the only infant food prepared according to the known laws of physiology.



**DOLIBER-GOODALE CO.,**

**BOSTON, MASS.**

18636  
THE ERA KEY TO THE U. S. P.

---

A COMPLETE LIST

OF THE

DRUGS AND PREPARATIONS

OF

The United States Pharmacopœia

REVISION OF 1890-3.

Giving Official Titles, Common Names and Synonyms  
of the Drugs, Chemicals and Preparations  
of the Pharmacopœia,

WITH DOSES

In Apothecaries Weights and Measures with  
Equivalents in Metric Terms.

---

COMPILED BY

THE PHARMACEUTICAL ERA,

D. O. HAYNES & CO., PUBLISHERS,

DETROIT, MICH,

1893.

**T**HE object of this little publication is to further the introduction and employment of the official drugs and preparations contained in our National standard, the United States Pharmacopœia.

As a companion to that invaluable work, we have endeavored to give the physicians and pharmacists of America, in convenient shape for everyday use and quick reference, such a "key" to the contents of that publication as will assist them in their work as prescribers and dispensers.

To all physicians and pharmacists who believe in the advancement of legitimate pharmacy, this little work is respectfully dedicated.

**THE PUBLISHERS.**



# THE ERA KEY TO THE U. S. P.

---

## DRUGS AND PREPARATIONS OF THE United States Pharmacopœia WITH DOSES.

---

### **Absinthium—*Absinthium*—Wormwood.**

The leaves and tops of *Artemisia Absinthium*.

DOSE: 20 to 40 grs. (1.3 to 2.6 gms.)

### **Acacia—*Acacia*—Gum Arabic.**

A gummy exudation from *Acacia Senegal*.

Preps: Mucilago *Acaciæ*. Pulvis *Cretæ Aromaticus*.

### **Acetanilidum—*Acetanilid*—Phenylacetamide.**

An acetyl derivative of aniline.

DOSE: 3 to 15 grs. (0.194 to 0.97 gms.)

### **Acetum Opii—*Vinegar of Opium*.**

10% Opium.

DOSE: 5 to 15 min. (0.31 to 0.92 c. c.)

### **Acetum Scillæ—*Vinegar of Squill*.**

10% Squill.

DOSE: 10 to 30 min. (0.62 to 1.85 c. c.)

### **Acidum Aceticum—*Acetic Acid*.**

36% Acetic Acid.

Prep: Acidum Aceticum Dilutum.

### **Acidum Aceticum Dilutum—*Diluted Acetic Acid*.**

6% Acetic Acid.

DOSE: 2 to 4 fl. drs. (7.39 to 14.79 c. c.)

### **Acidum Aceticum Glaciale—*Glacial Acetic Acid*.**

Absolute Acetic Acid.

### **Acidum Arsenosum—*Arsenous Acid*—Arsenic Trioxide. White Arsenic.**

Preps: Liquor Acidi Arsenosi. Liquor Potassii Arsenitis.

DOSE: 1-60 to 1-16 gr. (0.00108 to 0.00405 gms.)

---

**MELLIN'S FOOD**  
Is a true Liebig's Food.

**Acidum Benzoicum—Benzoic Acid.**

Obtained from Benzoin.

DOSE; 10 to 30 grs. (0.648 to 1.944 gms.)

**Acidum Boricum—Boric Acid—Boracic Acid.**

DOSE: 10 to 30 grs. (0.648 to 1.944 gms.)

Prep: Glyceritum Boroglycerini.

**Acidum Carbolicum—Carbolic Acid—Phenol.**

A constituent of coal tar.

DOSE: 1 to 3 grs. (0.0648 to 0.1944 gms.)

Prep: Glyceritum Acidi Carbolici.

**Acidum Carbolicum Crudum—Crude Carbolic Acid.**

From coal tar, chiefly cresol and phenol.

**Acidum Chromicum—Chromic Acid—Chromic Trioxide. Chromic Anhydride.**

DOSE: 1-24 to 1-4 gr. (0.0027 to 0.0162 gms.)

**Acidum Citricum—Citric Acid.**

Usually prepared from lemon juice.

DOSE: 10 to 30 grs. (0.648 to 1.944 gms.)

Prep: Syrupus Acidi Citrici.

**Acidum Gallicum—Gallic Acid.**

An organic acid, usually prepared from tannic acid.

DOSE: 5 to 30 grs. (0.324 to 1.944 gms.)

**Acidum Hydrobromicum Dilutum—Diluted Hydrobromic Acid.**

10% absolute Hydrobromic Acid.

DOSE:  $\frac{1}{2}$  to 2 fl. drs. (1.85 to 7.39 c. c.)

**Acidum Hydrochloricum—Hydrochloric Acid—Muriatic Acid.**

31.9% absolute Hydrochloric Acid.

DOSE; 5 to 10 min. diluted (0.31 to 0.62 c. c.)

Preps; Acidum Hydrochloricum Dilutum. Acidum Nitrohydrochloricum. Acidum Nitrohydrochloricum Dilutum.

**Acidum Hydrochloricum Dilutum—Diluted Hydrochloric Acid—Diluted Muriatic Acid.**

10% Hydrochloric Acid.

DOSE; 10 to 30 min. (0.62 to 1.85 c. c.)

**Acidum Hydrocyanicum Dilutum—Diluted Hydrocyanic Acid—Prussic Acid.**

2% absolute Hydrocyanic Acid.

DOSE: 1 to 3 min. (0.06 to 0.18 c. c.)

**Acidum Hypophosphorosum Dilutum—Diluted Hypophosphorous Acid.**

10% absolute Hypophosphorous Acid.

DOSE: 10 to 60 min. (0.62 to 3.70 c. c.)

---

**TRY MELLIN'S FOOD**  
For Hand-fed Infants.

**Acidum Lacticum—Lactic Acid.**

An organic acid, usually obtained by subjecting milk-sugar or grape-sugar to lactic fermentation.

75% absolute Lactic Acid.

DOSE:  $\frac{1}{2}$  to 2 fl. drs. (1.85 to 7.39 c. c.)

Prep: Syrupus Calcii Lactophosphatis.

**Acidum Nitricum—Nitric Acid.**

68% absolute Nitric Acid.

Preps: Acidum Nitricum Dilutum. Acidum Nitrohydrochloricum. Acidum Nitrohydrochloricum Dilutum.

**Acidum Nitricum Dilutum—Diluted Nitric Acid.**

Nitric Acid 10%.

DOSE: 5 to 30 min. (0.31 to 1.85 c. c.)

**Acidum Nitrohydrochloricum—Nitrohydrochloric Acid—Nitromuriatic Acid.**

Nitric Acid 18%. Hydrochloric Acid 82%.

DOSE: 2 to 5 min. (0.12 to 0.31 c. c.)

**Acidum Nitrohydrochloricum Dilutum—Diluted Nitrohydrochloric Acid—Diluted Nitromuriatic Acid.**

Nitric Acid 4%. Hydrochloric Acid 18%.

DOSE: 10 to 20 min. (0.62 to 1.24 c. c.)

**Acidum Oleicum—Oleic Acid.**

An organic acid obtained by cooling commercial Oleic Acid to 5 deg. C., and separating the liquid portion.

**Acidum Phosphoricum—Phosphoric Acid.**

85% absolute Orthophosphoric Acid.

DOSE: 3 to 7 min. (0.18 to 0.43 c. c.)

Prep: Acidum Phosphoricum Dilutum.

**Acidum Phosphoricum Dilutum—Diluted Phosphoric Acid.**

10% Phosphoric Acid.

DOSE: 5 to 30 min. (0.31 to 1.85 c. c.)

**Acidum Salicylicum—Salicylic Acid.**

From Carbolic Acid.

DOSE: 10 to 25 grs. (0.648 to 1.620 grs.)

**Acidum Stearicum—Stearic Acid.**

From solid fats, chiefly tallow.

**Acidum Sulphuricum—Sulphuric Acid.**

92.5% absolute Sulphuric Acid.

Preps: Acidum Sulphuricum Aromaticum. Acidum Sulphuricum Dilutum.

**Acidum Sulphuricum Aromaticum—Aromatic Sulphuric Acid.**

20% Sulphuric Acid.

DOSE: 5 to 10 min. (0.31 to 0.62 c. c.)

---

**TRY MELLIN'S FOOD**  
**For Invalids.**

**Acidum Sulphuricum Dilutum—Diluted Sulphuric Acid.**

10% Sulphuric Acid.

DOSE: 10 to 20 min. (0.62 to 1.23 c. c.)

**Acidum Sulphurosum—Sulphurous Acid.**

6.4% Sulphurous Acid Gas.

DOSE.  $\frac{1}{4}$  to 1 fl. dr (0.92 to 3.70 c. c.)**Acidum Tannicum—Tannic Acid—Gallotannic Acid. Digallic Acid.**

From Nutgall.

DOSE: 3 to 10 grs. (0.1944 to 0.648 gms.)

Preps: Colloidium Stypticum. Glyceritum Acidi Tannici.

Trochisci Acidi Tannici. Unguentum Acidi Tannici

**Acidum Tartaricum—Tartaric Acid.**

From argols.

DOSE: 10 to 30 grs. (0.648 to 1.944 gms.)

**Aconitum—Aconite.**The tuber of *Aconitum Napellus*.DOSE.  $\frac{1}{2}$  to 2 grs. (0.0324 to 0.129 gms.)

Preps: Extractum Aconiti. Extractum Aconiti Fluidum.

Tinctura Aconiti.

**Adeps—Lard.**Prepared internal fat of the abdomen of *Sus Scrofa*, Linne.

Preps: Adeps Benzoinatus. Ceratum (Compound Cerates).

Unguentum.

**Adeps Benzoinatus—Benzoinated Lard.****Adeps Lanæ Hydrosus—Hydrous Wool-fat.**

Purified fat of the wool of sheep.

**Æther—Ether—Æther Fortior, U. S. P., 1880.**

96% absolute Ether or Ethyl Oxide.

DOSE: 5 to 60 min. (0.31 to 3.70 c. c.)

Preps: Spiritus Ætheris. Spiritus Ætheris Comp.

**Æther Aceticus—Acetic Ether.**

98.5% of Ethyl Acetate.

DOSE 10 to 30 min. (0.62 to 1.86 c. c.)

**Alcohol—Alcohol.**

91% by weight or 94% by volume of Ethyl Alcohol.

Prep: Alcohol Dilutum.

**Alcohol Absolutum—Absolute Alcohol.**

Ethyl Alcohol, containing not more than 1% by weight of water.

**Alcohol Deodoratum—Deodorized Alcohol.**

92.5% by weight, or 95.1% by volume, of Ethyl Alcohol.

**Alcohol Dilutum—Diluted Alcohol.**

41% by weight, or 48.6% by volume, of Ethyl Alcohol.

**TRY MELLIN'S FOOD****In Typhoid Fever.**

**Allium—Garlic.**

The bulb of *Allium Sativum*.

DOSE: 15 to 60 grs. (0.972 to 3.888 gms.)

Prep: Syrupus Allii.

**Aloe Barbadosensis—Barbadoes Aloes.**

The inspissated juice of the leaves of *Aloe Vera*.

DOSE. 2 to 10 grs. (0.1296 to 0.648 gms.)

**Aloe Purificata—Purified Aloes.**

DOSE:  $\frac{1}{2}$  to 10 grs. (0.6324 to 0.648 gms.)

Preps: Extractum Colocynthis Compositum. Pilulæ Aloes. Pilulæ Aloes et Asafoetida. Pilulæ Aloes et Ferri. Pilulæ Aloes et Mastiches. Pilulæ Aloes et Myrrhae. Pilulæ Rhei Compositæ. Tinctura Aloes. Tinctura Aloes et Myrrhae. Tinctura Benzoini Composita.

**Aloe Socotrina—Socotrine Aloes.**

The inspissated juice of the leaves of *Aloe Perryi*.

DOSE 5 to 20 grs. (0.324 to 1.296 gms.)

Preps. Aloe Purificata. Extractum Aloes.

**Aloinum—Aloin.**

From Aloes.

DOSE.  $\frac{1}{4}$  to 2 grs. (0.0162 to 0.1296 gms.)

**Althæa—Althæa—Marshmallow.**

The root of *Althæa Officinalis*.

Prep: Syrupus Althææ.

**Alumen—Alum—Potassium Alum. Aluminum and Potassium Sulphate.**

DOSE: 10 to 60 grs. (0.648 to 3.888 gms.)

Prep: Alumen Exsiccatum.

**Alumen Exsiccatum—Dried Alum—Alumen Ustum. Burnt Alum.**

DOSE: 5 to 30 grs. (0.324 to 1.944 gms.)

**Alumini Hydras—Aluminum Hydrate—Aluminum Hydroxide. Hydrated Alumina.**

DOSE: 1 to 10 grs. (0.0648 to 0.648 gms.)

**Alumini Sulphas—Aluminum Sulphate.****Ammoniacum—Ammoniac.**

A gum resin obtained from *Dorema Ammoniacum*.

DOSE: 5 to 30 grs. (0.324 to 1.944 gms.)

Preps: Emplastrum Ammoniaci cum Hydrargyro. Emulsum Ammoniaci.

**Ammonii Benzoas—Ammonium Benzoate.**

DOSE: 10 to 30 grs. (0.648 to 1.944 gms.)

**Ammonii Bromidum—Ammonium Bromide.**

DOSE. 10 to 60 grs. (0.648 to 3.888 gms.)

**Ammonii Carbonas—Ammonium Carbonate.**

DOSE: 5 to 15 grs. (0.324 to 0.972 gms.)

Prep. Spiritus Ammoniae Aromaticus.

**TRY MELLIN'S FOOD**  
**In Cholera Infantum.**

**Ammonii Chloridum—Ammonium Chloride.**

DOSE; 5 to 30 grs. (0.324 to 1.944 gms.)

Prep: Trochisci Ammonii Chloridi.

**Ammonii Iodidum—Ammonium Iodide.**

DOSE. 3 to 10 grs. (0.194 to 0.648 gms.)

**Ammonii Nitras—Ammonium Nitrate.**

DOSE. 10 to 30 grs. (0.648 to 1.944 gms.)

**Ammonii Valerianas—Ammonium Valerianate.**

DOSE. 2 to 8 grs. (0.129 to 0.518 gms.)

**Amygdala Amara—Bitter Almond.**The seed of *Prunus Amygdalus* var. *Amara*.

Prep: Syrupus Amygdalæ.

**Amygdala Dulcis—Sweet Almond.**The seed of *Prunus Amygdalus* var. *Dulcis*.

Preps. Emulsum Amygdalæ. Syrupus Amygdalæ.

**Amyl Nitris—Amyl Nitrite.**

80% Amyl Nitrite.

DOSE: 2 to 4 min. (0.12 to 0.25 c. c.)

**Amylum—Starch.**The fecula of the seed of *Zea Mays*.

Prep; Glyceritum Amyli.

**Anisum—Anise.**The fruit of *Pimpinella Anisum*.

DOSE: 10 to 30 grs. (0.648 to 1.944 gms.)

**Anthemis—Anthemis—Chamomile.**The flower heads of *Anthemis Nobilis*.DOSE:  $\frac{1}{2}$  to 2 drs. (1.944 to 7.776 gms.)**Antimonii et Potassii Tartras—Antimony and Potassium Tartrate—Tartar Emetic, Tartarated Antimony.**DOSE.  $\frac{1}{2}$  to 3 grs. (0.0081 to 0.194 gms.)

Preps Syrupus Scillæ Compositus. Vinum Antimonii.

**Antimonii Oxidum—Antimony Oxide—Antimony Trioxide.**

DOSE 1 to 3 grs. (0.0648 to 0.194 gms.)

Prep. Pulvis Antimonialis.

**Antimonii Sulphidum—Antimony Sulphide—Antimony Trisulphide.**

Prep: Antimonii Sulphidum Purificatum.

**Antimonii Sulphidum Purificatum—Purified Antimony Sulphide—Purified Antimony Trisulphide.**

Prep: Antimonium Sulphuratum.

---

**TRY MELLIN'S FOOD**  
In Intestinal Indigestion.

**Antimonium Sulphuratum** — *Sulphurated Antimony*—Kermes Mineral.

DOSE: 1 to 3 grs. (0.0648 to 0.194 gms.)

Prep: *Pilulæ Antimonii Compositæ*.

**Apocynum**—*Apocynum*—Canadian Hemp.

The root of *Apocynum Cannabinum*.

DOSE: 2 to 20 grs. (0.1296 to 1.296 gms.)

Prep: *Extractum Apocyni Fluidum*.

**Apomorphinæ Hydrochloras**—*Apomorphine Hydrochlorate*.

Hydrochlorate of an artificial alkaloid, prepared from Morphine or Codeine.

DOSE: 1-16 to  $\frac{1}{4}$  gr. (0.004 to 0.0081 gm.)

**Aqua**—*Water*.

Prep: *Aqua Destillata*.

**Aqua Ammoniæ**—*Ammonia Water*.

An aqueous solution of Ammonia, containing 10% by weight of the gas.

DOSE: 10 to 30 min. (0.62 to 1.85 c. c.)

Prep: *Linimentum Ammoniæ*. *Spiritus Ammoniæ Aromaticus*.

**Aqua Ammoniæ Fortior**—*Stronger Ammonia Water*.

An aqueous solution of Ammonia, containing 28% of the gas.

Prep: *Spiritus Ammoniæ*.

**Aqua Amygdalæ Amaræ**—*Bitter Almond Water*.

DOSE:  $\frac{1}{2}$  to 2 fl. drs. (1.85 to 7.39 c. c.)

**Aqua Anisi**—*Anise Water*.

DOSE:  $\frac{1}{4}$  to 1 fl. oz. (7.39 to 29.57 c. c.)

**Aqua Aurantii Florum**—*Orange Flower Water*.

Prep: *Syrupus Aurantii Florum*.

**Aqua Aurantii Florum Fortior**—*Stronger Orange Flower Water*—Triple Orange Flower Water.

Prep: *Aqua Aurantii Florum*.

**Aqua Camphoræ**—*Camphor Water*.

Camphor 0.8%.

DOSE:  $\frac{1}{2}$  to 2 fl. ozs. (14.79 to 59.10 c. c.)

**Aqua Chlorig**—*Chlorine Water*.

An aqueous solution of Chlorine, containing at least .4% of the gas.

DOSE: 1 to 2 fl. ozs. (3.70 to 14.79 c. c.)

---

**TRY MELLIN'S FOOD**  
As a diet for Nursing Mothers.

**Aqua Chloroformi—Chloroform Water.**

DOSE: 1 to 4 fl. drs. (3.70 to 14.79 c. c.)

**Aqua Cinnamomi—Cinnamon Water.**DOSE:  $\frac{1}{2}$  to 1 fl. oz. (14.79 to 29.57 c. c.)**Aqua Creosoti—Creosote Water.**

DOSE: 2 to 4 fl. drs. (7.39 to 14.79 c. c.)

**Aqua Destillata—Distilled Water.****Aqua Fœniculi—Fennel Water.**DOSE:  $\frac{1}{4}$  to 1 fl. oz. (7.39 to 29.57 c. c.)**Aqua Hydrogenii Dioxidii—Solution of Hydrogen Dioxide—Solution of Hydrogen Peroxide.**

A slightly acid, aqueous solution of Hydrogen Dioxide, containing, when freshly prepared, about 3% by weight of the pure Dioxide, corresponding to about ten volumes of available oxygen.

**Aqua Menthæ Piperitæ — Peppermint Water.**DOSE:  $\frac{1}{2}$  to 2 fl. ozs. (14.79 to 59.10 c. c.)**Aqua Menthæ Viridis—Spearmint Water.**DOSE:  $\frac{1}{2}$  to 2 fl. ozs. (14.79 to 59.10 c. c.)**Aqua Rosæ—Rose Water.**

Prep: Unguentum Aquæ Rosæ.

**Aqua Rosæ Fortior—Stronger Rose Water—Triple Rose Water.**

Water saturated with the volatile oil of rose petals, obtained as a by-product in the distillation of Oil of Rose.

Prep: Aqua Rosæ.

**Argenti Cyanidum—Silver Cyanide.**

DOSE: 1.60 to 1.20 gr. (0.0010 to 0.0032 gms.)

Prep: Acidum Hydrocyanicum Dilutum.

**Argenti Iodidum—Silver Iodide.**DOSE:  $\frac{1}{2}$  to 2 grs. (0.0324 to 0.1296 gms.)**Argenti Nitras—Silver Nitrate.**DOSE:  $\frac{1}{4}$  to  $\frac{1}{2}$  gr. (0.0081 to 0.0324 gms.)

Prep: Argenti Nitras Dilutus. Argenti Nitras Fusus.

**Argenti Nitras Dilutus—Diluted Silver Nitrate—Mitigated Caustic.****Argenti Nitras Fusus—Moulded Silver Nitrate—Lunar Caustic.****Argenti Oxidum—Silver Oxide.**DOSE:  $\frac{1}{2}$  to 2 grs. (0.0324 to 0.1296 gms.)**Arnica Flores—Arnica Flowers.**

The flower heads of Arnica Montana.

DOSE: 10 to 20 grs. (0.648 to 1.296 gms.)

Prep: Tinctura Arnicae Florum.

**MELLIN'S FOOD****Makes cow's milk easily digestible.**



**Arnica Radix—Arnica Root.**

The rhizome and roots of *Arnica Montana*.

DOSE: 5 to 20 grs. (0.324 to 1.296 gms.)

Preps: Extractum Arnica Radicis. Extractum Arnica Radicis Fluidum. Tinctura Arnica Radicis.

**Arseni Iodidum—Arsenic Iodide.**

DOSE: 1-24 to  $\frac{1}{4}$  gr. (0.0027 to 0.0081 gms.)

Prep: Liquor Arseni Hydrargyri Iodidi.

**Asafoetida—Asafoetida.**

A gum-resin obtained from the root of *Ferula Nanthex* and of *Ferula Foetida*.

DOSE: 5 to 15 grs. (0.324 to 0.972 gms.)

Preps: Emulsum Asafoetidae. Pilulae Aloes et Asafoetidae. Pilulae Asafoetidae. Tinctura Asafoetidae.

**Asclepias—Asclepias.**

The root of *Asclepias Tuberosa*.

DOSE:  $\frac{1}{2}$  to 2 drs. (1.944 to 7.776 gms.)

Prep: Extractum Asclepiadis Fluidum.

**Aspidium—Aspidium—Male Fern.**

The rhizome of *Dryopteris Filix-mas* and of *Dryopteris Marginalis*.

DOSE:  $\frac{1}{2}$  to 1 $\frac{1}{2}$  drs. (1.944 to 5.932.)

Prep: Oleoresina Aspidii.

**Aspidosperma—Aspidosperma—Quebracho.**

The bark of *Aspidosperma Quebracho-blanco*.

DOSE: 5 to 30 grs. (0.324 to 1.944 gms.)

Prep: Extractum Aspidospermatis Fluidum.

**Atropina—Atropine.**

An alkaloid, obtained from *Belladonna*.

DOSE: 1-96 to 1-64 gr. (0.0006 to 0.001 gm.)

**Atropinae Sulphas—Atropine Sulphate.**

DOSE: 1-96 to 1-48 gr. (0.0006 to 0.0013 gm.)

**Aurantii Amari Cortex—Bitter Orange Peel.**

The rind of the fruit of *Citrus vulgaris*.

Preps. Extractum Aurantii Amari Fluidum. Tinctura Aurantii Amari.

**Aurantii Dulcis Cortex—Sweet Orange Peel.**

The rind of the fresh fruit of *Citrus Aurantium*.

Preps: Syrupus Aurantii. Tinctura Aurantii Dulcis.

**Aurii et Sodii Chloridum—Gold and Sodium Chloride.**

A mixture of equal parts by weight of dry Gold Chloride and Sodium Chloride.

DOSE: 1-16 to 1-10 gr. (0.004 to 0.0065 gm.)

**Balsamum Peruvianum—Balsam of Peru.**

A Balsam obtained from *Tolulfera Peruviana*.

DOSE: 10 to 30 min. (0.62 to 1.85 c. c.)

---

**MELLIN'S FOOD**  
Is readily assimilated.

**Balsamum Tolutanum—Balsam of Tolu.**

A balsam obtained from *Tolulfera Balsamum*.

DOSE: 10 to 30 min. (0.62 to 1.85 c. c.)

Preps: *Syrupus Tolutanus*. *Tinctura Benzoini Composita*  
*Tinctura Tolutana*.

**Barii Dioxidum—Barium Dioxide—Barium Peroxide.**

Commercial, anhydrous Barium Dioxide.

Prep: *Aqua Hydrogenii Dioxidi*.

**Belladonnæ Folia—Belladonna Leaves.**

The leaves of *Atropa Belladonna*.

DOSE: 1 to 5 grs. (0.0648 to 0.324 gm.)

Preps: *Extractum Belladonnæ Foliorum Alcoholicum*.  
*Tinctura Belladonnæ Foliorum*.

**Belladonnæ Radix—Belladonna Root.**

The root of *Atropa Belladonna*.

DOSE: 1 to 3 grs. (0.0648 to 0.1944 gm.)

Prep: *Extractum Belladonnæ Radicis Fluidum*.

**Benzinum—Benzin—Petroleum Benzin. Petroleum Ether.**

A purified distillate from American petroleum, consisting of hydrocarbons chiefly of the marsh-gas series and homologous compounds.

**Benzoinum—Benzoin.**

A balsamic resin obtained from *Styrax Benzoin*.

Preps: *Adeps Benzoinatus*. *Tinctura Benzoini*. *Tinctura Benzoini Composita*.

**Bismuthi Citras—Bismuth Citrate.**

DOSE: 1 to 5 grs. (0.0648 to 0.324 gm.)

Prep: *Bismuthi et Ammonii Citras*.

**Bismuthi et Ammonii Citras—Bismuth and Ammonium Citrate.**

DOSE: 1 to 5 grs. (0.0648 to 0.324 gm.)

**Bismuthi Subcarbonas—Bismuth Subcarbonate.**

DOSE: 10 to 60 grs. (0.648 to 3.888 gms.)

**Bismuthi Subnitras—Bismuth Subnitrate.**

DOSE: 10 to 60 grs. (0.648 to 3.888 gms.)

**Bromum—Bromine.****Bryonia—Bryonia—Bryony.**

The root of *Bryonia alba* and of *Bryonia dioica*.

DOSE: 10 to 60 grs. (0.648 to 3.888 gms.)

Prep: *Tinctura Bryoniæ*.

**Buchu—Buchu.**

The leaves of *Barosma betulina* and *Barosma crenulata*.

DOSE: 15 to 30 grs. (0.972 to 1.944 gms.)

Prep: *Extractum Buchu Fluidum*.

**MELLIN'S FOOD**

Is a soluble dry extract from wheat and malt.

**Caffeina—Caffeine—Theine.**

A feebly basic principle from the dried leaves of *Thea sinensis* or from the dried seeds of *Coffea arabica*, and found also in other plants.

DOSE: 1 to 3 grs. (0.0648 to 0.1944 gms.)

Preps: *Caffeina Citrata*. *Caffeina Citrata Effervescens*.

**Caffeina Citrata—Citrated Caffeine.**

DOSE: 2 to 5 grs. (0.1296 to 0.324 gms.)

**Caffeina Citrata Effervescens—Effervescent Citrated Caffeine.**

DOSE: 1 to 3 drs. (3.888 to 11.664 gms.)

**Calamus—Calamus—Sweet Flag.**

Rhizome of *Acorus Calamus*.

DOSE. *Ad libitum*.

Prep: *Extractum Calami Fluidum*.

**Calci Bromidum—Calcium Bromide.**

DOSE. 10 to 60 grs. (0.648 to 3.888 gms.)

**Calci Carbonas Præcipitatus—Precipitated Calcium Carbonate.**

DOSE: 5 to 60 grs. (0.324 to 3.888.)

**Calci Chloridum—Calcium Chloride.**

Calcium Chloride rendered anhydrous by fusion at the lowest possible temperature.

DOSE: 5 to 20 grs. (0.324 to 1.296 gms.)

**Calci Hypophosphis—Calcium Hypophosphate.**

DOSE: 5 to 30 grs. (0.324 to 1.944 gms.)

Prep: *Syrupus Hypophosphitum*.

**Calci Phosphas Præcipitatus—Precipitated Calcium Phosphate.**

DOSE: 5 to 30 grs. (0.324 to 1.944 gms.)

Prep: *Syrupus Calci Lactophosphatis*.

**Calci Sulphas Exsiccatus—Dried Calcium Sulphate.****Calendula—Calendula—Marigold.**

The florets of *Calendula officinalis*.

DOSE: 15 to 60 grs. (0.972 to 3.888 gms.)

Prep: *Tinctura Calendulæ*.

**Calumba—Calumba—Columbo.**

The root of *Jateorhiza palmata*.

DOSE. 5 to 30 grs. (0.324 to 1.944 gms.)

Preps: *Extractum Calumbæ Fluidum*. *Tinctura Calumbæ*.

**MELLIN'S FOOD**

Is entirely free from starch and cane sugar.

**Calx—Lime.**

Lime prepared by burning white marble, oyster shells, or the purest varieties of calcium carbonate.

Preps: Liquor Calcis. Potassa cum Calce. Syrupus Calcis.

**Calx Chlorata—Chlorinated Lime.**

Containing not less than 35% of available chlorine.

DOSE: 3 to 6 grs. (0.0194 to 0.388 gms.)

**Calx Sulphurata—Sulphurated Lime—Crude Calcium Sulphide.**

DOSE. 1-10 to 1 gr. (0.0064 to 0.0648 gms.)

**Cambogia—Gamboge.**

A gum-resin obtained from *Garcinia Hanburii*.

DOSE: 1 to 5 grs. (0.0648 to 0.324 gms.)

Prep: Pilulæ Catharticæ Compositæ.

**Camphora—Camphor.**

A stearopten (having the nature of a ketone) obtained from *Cinnamomum Camphora* and purified by sublimation.

DOSE: 1 to 10 grs. (0.0648 to 0.648 gms.)

Preps: Aqua Camphoræ. Linimentum Camphoræ. Linimentum Saponis. Spiritus Camphoræ. Tinctura Opii Camphorata.

**Camphora Monobromata—Monobromated Camphor.**

DOSE. 1 to 5 grs. (0.0648 to 0.324 gms.)

**Cannabis Indica—Indian Cannabis—Indian Hemp.**

The flowering tops of the plant of *Cannabis sativa*.

DOSE: 2 to 5 grs. (0.1296 to 0.324 gms.)

Preps: Extractum Cannabis Indicæ. Extractum Cannabis Indicæ Fluidum. Tinctura Cannabis Indicæ.

**Cantharis—Cantharides—Spanish Flies.**

*Cantharis vesicatoria*.

DOSE:  $\frac{1}{2}$  to 2 grs. (0.0324 to 0.1296 gms.)

Preps: Ceratum Cantharidis. Collodium Cantharidatum. Tinctura Cantharidis.

**Capsicum—Capsicum—Cayenne Pepper. African Pepper.**

The fruit of *Capsicum fastigiatum*.

DOSE: 3 to 5 grs. (0.1944 to 0.324 gms.)

Preps: Extractum Capsici Fluidum. Oleoresina Capsici. Tinctura Capsici.

**Carbo Animalis—Animal Charcoal.**

Charcoal prepared from bone.

Prep: Carbo Animalis Purificatus.

---

**MELLIN'S FOOD**

Is unrivalled as an artificial food for children.

**Carbo Animalis Purificatus**—*Purified Animal Charcoal.*

**Carbo Ligni**—*Charcoal.*

Charcoal prepared from soft wood and very finely powdered.

DOSE: 1 to 4 grs. (0.0648 to 0.2592 gms.)

**Carbonei Disulphidum**—*Carbon Disulphide.*  
(Carbonei Bisulphidum, U. S. P., 1880.)

DOSE. 5 to 10 drops (0.31 to 0.62 c. c.)

**Cardamomum**—*Cardamom.*

The fruit of *Elletaria repens*.

DOSE 10 to 15 grs. (0.648 to 0.972 gms.)

Preps Pulvis Aromaticus. Tinctura Cardamomi. Tinctura Cardamomi Composita.

**Carum**—*Caraway.*

The fruit of *Carum Carvi*.

DOSE. 15 to 30 grs. (0.972 to 1.944 gms.)

Prep. Tinctura Cardamomi Composita.

**Caryophyllus**—*Cloves.*

The unexpanded flowers of *Eugenia Aromatica*.

DOSE 5 to 30 grs. (0.324 to 1.944 gms.)

Prep. Tinctura Lavandulae Composita.

**Cascarilla**—*Cascarilla.*

The bark of *Croton Eluteria*.

DOSE 10 to 30 grs. (0.648 to 1.944 gms.)

**Cassia Fistula**—*Cassia Fistula*—*Purging Cassia.*

The fruit of *Cassia Fistula*.

DOSE 1 to 8 drs. (3.888 to 31.103 gms.)

Prep. Confectio Sennæ.

**Castanea**—*Castanea*—*Chestnut.*

The leaves of *Castanea dentata*, collected in September or October while it is green.

DOSE:  $\frac{1}{2}$  to 2 drs. (1.944 to 7.776 gms.)

Prep. Extractum Castaneæ Fluidum.

**Catechu**—*Catechu*

An extract prepared from the wood of *Acacia Catechu*.

DOSE 10 to 30 grs. (0.648 to 1.944 gms.)

Preps Tinctura Catechu Composita. Trochisci Catechu.

**Caulophyllum**—*Caulophyllum*—*Blue Cohosh.*

The rhizome and roots of *Caulophyllum thalictroides*.

DOSE 5 to 30 grs. (0.324 to 1.944 gms.)

**Cera Alba**—*White Wax.*

Yellow wax bleached.

Prep: Ceratum (Compound Cerates).

## MELLIN'S FOOD

Stimulates the bowels to regular action.

**Cera Flava—Yellow Wax.**

A peculiar, concrete substance, prepared by *Apis mellifica*.  
 Preps. *Ceratum Resinæ*. (Compound Cerates.) *Emplastrum Resinæ Unguentum*.

**Ceratum—Cerate.****Ceratum Camphoræ—Camphor Cerate.****Ceratum Cantharidis—Cantharides Cerate.**

*Cantharides* 32%.

Prep; *Emplastrum Picis Cantharidatum*.

**Ceratum Cetacei—Spermaceti Cerate.**

*Spermaceti* 10%.

**Ceratum Plumbi Subacetatis—Cerate of Lead Subacetate—Goulard's Cerate.**

Solution of Lead Subacetate 20%.

**Ceratum Resinæ—Resin Cerate—Basilicon Ointment.**

Resin 35%.

**Ceril Oxalas—Cerium Oxalate—Cerous Oxalate.**

DOSE: 1 to 5 grs. (0.0648 to 0.324 gms.)

**Cetaceum—Spermaceti.**

A peculiar concrete fatty substance, obtained from *Physeter macrocephalus*.

Preps. *Ceratum Cetacei*. *Unguentum Aquæ Rosæ*.

**Cetraria—Cetraria—Iceland Moss.**

*Cetraria Islandica*.

Prep. *Decoctum Cetrariæ*.

**Charta Potassii Nitratis—Potassium Nitrate Paper.****Charta Sinapis—Mustard Paper.****Chelidonium—Chelidonium—Celandine.**

*Chelidonium majus*.

DOSE: 15 to 60 grs. (0.972 to 3.888 gms.)

**Chenopodium — Chenopodium — Wormseed.**

The fruit of *Chenopodium anthelminticum*.

DOSE: 15 to 30 grs. (0.972 to 1.944 gms.)

**Chimaphila—Chimaphila—Pipsissewa.**

Leaves of *Chimaphila umbellata*.

DOSE:  $\frac{1}{2}$  to 2 drs. (1.944 to 7.776 gms.)

Prep; *Extractum Chimaphilæ Fluidum*.

**Chirata—Chirata.**

*Swertia Chirata*.

DOSE: 20 grs. (1.296 gms.)

Preps. *Extractum Chiratæ Fluidum*. *Tinctura Chir*

**MELLIN'S FOOD**

Actually assists to digest milk or other food  
 with which it is mixed.

**Chloral—Chloral—Chloral Hydrate.**

A crystalline solid, composed of trichloraldehyde or chloral with one molecule of water.

DOSE: 10 to 45 grs. (0.648 to 2.916 gms.)

**Chloroformum—Chloroform—Chloroformum Purificatum. Pharm., 1880.**

A liquid containing at least 99 to 99.4% by weight of absolute Chloroform, and 1 to 0.6% of alcohol.

DOSE: 2 to 5 min. (0.18 to 0.31 c. c.)

Preps: Aqua Chloroformi. Emulsum Chloroformi. Linimentum Chloroformi. Spiritus Chloroformi.

**Chondrus—Chondrus—Irish Moss. Carragheen.**

*Chondrus Crispus* and *Gigartina mamillosa*.

**Chrysarobinum—Chrysarobin.**

A neutral principle, in its more or less commercial, impure form, extracted from Goa Powder, a substance found deposited in the wood of *Andira Araroba*.

DOSE: 1.64 to 1.12 gr. (0.0010 to 0.0054 gms.)

Prep: Unguentum Chrysarobini.

**Cimicifuga—Cimicifuga—Black Snakeroot.**

The rhizome and roots of *Cimicifuga racemosa*.

DOSE: 20 to 30 grs. (1.296 to 1.944 gms.)

Preps: Extractum Cimicifugæ. Extractum Cimicifugæ Fluidum. Tinctura Cimicifugæ.

**Cinchona—Cinchona.**

The bark of *Cinchona Callaya*, *Cinchona officinalis* and of hybrids of these and other species of *Cinchona*, containing not less than 5% of total alkaloids and at least 2.5% of quinine.

DOSE:  $\frac{1}{2}$  to 1 oz. (15.552 to 31.103 gms.)

Preps: Extractum Cinchonæ. Extractum Cinchonæ Fluidum. Infusum Cinchonæ. Tinctura Cinchonæ.

**Cinchona Rubra—Red Cinchona.**

The bark of *Cinchona succirubra*, containing not less than 5% of its peculiar alkaloids.

DOSE:  $\frac{1}{2}$  to 1 oz. (15.552 to 31.103 gms.)

Prep: Tinctura Cinchonæ Composita.

**Cinchonidinæ Sulphas—Cinchonidine Sulphate.**

The neutral sulphate of an alkaloid extracted from the bark of various species of *Cinchona*.

DOSE: 3 to 30 grs. (0.1944 to 1.944 gms.)

**Cinchonina—Cinchonine.**

An alkaloid obtained from the bark of various species of *Cinchona*.

DOSE: 1 to 30 grs. (0.0648 to 1.944 gms.)

---

**MELLIN'S FOOD**

Contains the nitrogenous and phosphatic matters essential for the growing child.

**Cinchoninæ Sulphas—Cinchonine Sulphate.**

DOSE. 1 to 30 grs. (0.0648 to 1.944 gms.)

**Cinnamomum Cassia—Cassia Cinnamon—Cassia Bark.**The bark of the shoots of one or more undetermined species of *Cinnamomum* grown in China (Chinese Cinnamon).

DOSE: 10 to 30 grs. (0.648 to 1.944 gms.)

Preps: *Tinctura Cardamomi Composita*. *Tinctura Catechu Composita*. *Tinctura Lavandulæ Composita*.**Cinnamomum Salgonicum—Saigon Cinnamon.**The bark of an undetermined species of *Cinnamomum*.

DOSE: 10 to 30 grs. (0.648 to 1.944 gms.)

**Cinnamomum Zeylanicum—Ceylon Cinnamon.**The inner bark of the shoots of *Cinnamomum zeylanicum*.

DOSE: 10 to 30 grs. (0.648 to 1.944 )

Preps: *Pulvis Aromaticus*. *Tinctura Cinnamomi*.**Coca—Erythroxylon.**The leaves of *Erythroxylon Coca*.DOSE:  $\frac{1}{2}$  to 1 dr. (1.944 to 3.888 gms.)Prep: *Extractum Cocæ Fluidum*.**Cocainæ Hydrochloras—Cocaine Hydrochlorate.**

The hydrochlorate of an alkaloid extracted from Coca.

DOSE.  $\frac{1}{4}$  to 2 grs. (0.0081 to 0.1296 gms.)**Coccus—Cochineal.**The dried female of *Coccus cacti*.DOSE:  $\frac{1}{4}$  to  $\frac{1}{2}$  gr. (0.0108 to 0.0324 gms.)**Codeina—Codeine.**

An alkaloid extracted from Opium.

DOSE:  $\frac{1}{4}$  to 3 grs. (0.0162 to 0.1944 gms.)**Colchici Radix—Colchicum Root.**The corm of *Colchicum autumnale*.

DOSE: 2 to 8 grs. (0.1296 to 0.5184 gms.)

Preps: *Extractum Colchici Radicis*. *Extractum Colchici Radicis Fluidum*. *Vinum Colchici Radicis*.**Colchici Semen—Colchicum Seed.**The seed of *Colchicum autumnale*.

DOSE: 1 to 5 grs. (0.0648 to 0.324 gms.)

Preps: *Extractum Colchici Seminis Fluidum*. *Tinctura Colchici Seminis*. *Vinum Colchici Seminis*.**Collodium—Collodion.**Preps: *Collodium Flexile*. *Collodium Stypticum*.**MELLIN'S FOOD**Prepared with milk fulfils every requirement  
of a food for infan's.



**Collodium Cantharidatum — Cantharidal**

### Collodion—Blistering Collodion.

Cantharides 60%.

**Collodium Flexile**—*Flexible Collodion.*

**Collodium Stypticum**—*Styptic Collodion.*

**Colocynthis**—*Colocynth.*

The fruit of *Citrullus Colocynthis* deprived of its rind.

**DOSE.** 2 to 8 grs. (0.1296 to 0.5184 gms.)

Prep    *Extractum Colocynthis.*

**Confectio Rosæ**—*Confection of Rose.*

**Confectio Sennæ**—*Confection of Senna.*

**DOSE:** 2 to 4 drs. (7.776 to 15.552 gms.)

**Conium**—*Conium*—Hemlock.

The full grown fruit of *Conium maculatum*, gathered while yet green.

**DOSE:** 2 to 5 grs. (0.1296 to 0.324 gms.)

Preps. Extractum Conii

**Extractum Conii**

Fluidum.

**Convallaria—Convallaria.**

The rhizome and roots of *Convallaria majalis*.

**DOSE.** 5 to 30 grs. (0.324 to 1.944 gms.)

Prep. Extractum Convallariæ Fluidum.

**Copaiba**—*Copaiba*—Balsam of Copaiba.

The Oleoresin of *Copaiba Langsdorffii* and other species of *Copaifera*.

**DOSE.**  $\frac{1}{2}$  to 1 dr. (0.972 to 3.888 gms.)

Prep. Massa Copaibæ.

**Coriandrum—Coriander.**

The fruit of *Coriandrum sativum*.

**DOSE.** 10 to 30 grs. (0.648 to 1.944 gms.)

**Prep. Confectio Sennae.**

**Creosotum**—*Creosote*.

A mixture of phenols, chiefly guaiacol and creosol, obtained during the distillation of wood tar, preferably of that derived from the beech, *Fagus silvatica*.

**DOSE** 1 to 5 grs (0.0648 to 0.324 gms.)

Prep: Aqua Creosoti.

**Creta Præparata—Prepared Chalk.**

**DOSE** 10 to 60 grs. (0.648 to 3.888 gms.)

Preps: Hydrargyrum cum Creta. Pulvis Cretæ Compos-  
itus. Trochisci Cretæ.

**Crocus—Saffron.**

The stigmas of *Crocus sativus*.

**DOSE:** 5 to 30 grs. (0.324 to 1.944 gms.)

Prep: Tinctura Croci.

## MELLIN'S FOOD

Contains a large amount of proteids and soluble phosphates.

**Cubeba—Cubeb.**

The unripe fruit of *Piper Cubeba*.

DOSE:  $\frac{1}{4}$  to 3 drs. (0.972 to 11.664 gms.)

Preps: Extractum Cubebæ Fluidum. Oleoresina Cubebæ. Tinctura Cubebæ.

**Cupri Sulphas—Copper Sulphate—Cupric Sulphate.**

DOSE:  $\frac{1}{2}$  to 10 grs. (0.0324 to 0.648 gms.)

**Cusso - Kouso--Brayera.**

The female inflorescence of *Hagenia abyssinica*.

DOSE: 2 to 4 drs. (7.776 to 15.552 gms.)

Prep. Extractum Cusso Fluidum.

**Cypripedium - Cypriped'um—Ladies' Slipper.**

The rhizome and root of *Cypripedium pubescens* and of *Cypripedium parviflorum*.

DOSE: 15 to 30 grs. (0.972 to 1.944 gms.)

Prep: Extractum Cypripedi Fluidum.

**Decocta—Decoctions.**

An ordinary decoction, the strength of which is not directed by the physician, nor specified by the Pharmacopœia, shall be prepared by the following formula

The substance, coarsely comminuted, 50 gms.

Water, a sufficient quantity to make 1000 c. c.

**Decoctum Cetrariæ—Decoction of Cetraria.****Decoctum Sarsaparillæ Compositum—**

*Compound Decoction of Sarsaparilla.*

DOSE. 4 to 6 fl. ozs. (118.3 to 177.4 c. c.)

**Digitalis—Digitalis—Foxglove.**

The leaves of *Digitalis purpurea*, collected from plants of the second year's growth.

DOSE:  $\frac{1}{2}$  to 3 grs. (0.0324 to 0.1944 gms.)

Preps: Extractum Digitalis. Extractum Digitalis Fluidum. Infusum Digitalis. Tinctura Digitalis.

**Dulcamara—Dulcamara—Bittersweet.**

The young branches of *Solanum Dulcamara*.

DOSE: 1 to 2 drs. (3.888 to 7.776 gms.)

Prep: Extractum Dulcamaræ Fluidum.

**Elastica—Caoutchouc—India Rubber.**

The prepared milk juice of various species of *Hevea*, known in commerce as Para Rubber.

**Elaterinum—Elaterin.**

A neutral principle extracted from *Elaterium*, a substance deposited by the juice of the fruit of *Echallium Elaterium*.

DOSE. 1-32 to 1-16 gr. (0.0020 to 0.0040 gms.)

Prep: Trituratio Elaterini.

**Elixir Aromaticum—Aromatic Elixir.**

DOSE 1 to 3 fl. drs. (3.7 to 11.09 c. c.)

**MELLIN'S FOOD**

Is a most useful addition to the diet of healthy children.

**Elixir Phosphori—Elixir of Phosphorus.**

DOSE. 15 to 75 min. (0.92 to 4.62 c. c.)

**Emplastrum Ammoniaci cum Hydrargyro—Ammoniac Plaster with Mercury.****Emplastrum Arnicæ—Arnica Plaster.**

Extract of Arnica root, 33%.

**Emplastrum Belladonnæ—Belladonna Plaster.**

Alcoholic Extract of Belladonna, 20%.

**Emplastrum Capsici—Capsicum Plaster.****Emplastrum Ferri—Iron Plaster—Strengthening Plaster.**

Ferric Hydrate, 9%.

**Emplastrum Hydrargyri—Mercurial Plaster.**

Mercury, 30%. Oleate of Mercury, 12%.

**Emplastrum Ichthyocollæ—Isinglass Plaster—Court Plaster.****Emplastrum Opii—Opium Plaster**

Extract of Opium, 6%.

**Emplastrum Picis Burgundicæ—Burgundy Pitch Plaster.**

Burgundy Pitch, 80%.

**Emplastrum Picis Cantharidatum—Cantharidal Pitch Plaster—Warming Plaster.**

Cerate of Cantharides, 8%.

**Emplastrum Plumbi—Lead Plaster—Diachylon Plaster.**

Preps: Emplastrum Resinæ. Emplastrum Saponis. Unguentum Diachylon.

**Emplastrum Resinæ—Resin Plaster—Adhesive Plaster.**

Resin, 14%.

**Emplastrum Saponis—Soap Plaster.**

Soap, 10%.

**Emulsum Ammoniaci—Emulsion of Ammoniac. Mistura Ammoniaci, U. S. P., 1880.**

Ammoniac, 4%.

DOSE.  $\frac{1}{2}$  to 1 fl. oz. (14.79 to 29.57 c. c.)**Emulsum Amygdalæ—Emulsion of Almond—Mistura Amygdalæ, U. S. P., 1880. Milk of Almonds.**

Sweet Almond, 6%.

DOSE: 2 to 8 fl. ozs. (59.1 to 236.6 c. c.)

**MELLIN'S FOOD**

Has proved its greater nutritive value over all other artificial foods.

**Emulsum Asafoetidae**—*Emulsion of Asafoetida*  
—*Mistura Asafoetida*, U. S. P., 1880. Milk of  
*Asafoetida*.

*Asafoetida*, 4%.

DOSE: 2 to 4 fl. drs. (7.39 to 14.79 c. c.)

**Emulsum Chloroformi**—*Emulsion of Chloroform*—*Mistura Chloroformi*, U. S. P., 1880.

DOSE: 1 to 2 drs. (3.888 to 7.776 gms.)

**Ergota**—*Ergot*—Ergot of Rye.

The sclerotium of *Claviceps purpurea*, replacing the grain of rye. *Secale cereale*.

DOSE: 15 to 30 grs. (0.972 to 1.944 gms.)

Preps: *Extractum Ergotæ Fluidum*. *Vinum Ergotæ*.

**Eriodictyon**—*Eriodictyon*.

The leaves of *Eriodictyon glutinosum*.

DOSE: 15 to 30 grs. (0.972 to 1.944 gms.)

Prep: *Extractum Eriodictyi Fluidum*.

**Eucalyptol**—*Eucalyptol*.

A neutral body obtained from the volatile oil of *Eucalyptus globulus* and of some other species of *Eucalyptus*.

DOSE: 5 to 30 min. (0.31 to 1.85 c. c.)

**Eucalyptus**—*Eucalyptus*.

The leaves of *Eucalyptus globulus*, collected from the older parts of the tree.

DOSE:  $\frac{1}{2}$  to 2 drs. (1.944 to 7.776 gms.)

Prep: *Extractum Eucalypti Fluidum*.

**Euonymus**—*Euonymus*—Wahoo.

The bark of the root of *Euonymus atropurpureus*.

DOSE: 1 to 2 drs. (3.888 to 7.776 gms.)

Prep: *Extractum Euonymi*.

**Eupatorium**—*Eupatorium*—Thoroughwort.

The leaves and flowering tops of *Eupatorium perfoliatum*.

DOSE:  $\frac{1}{4}$  to 1 dr. (0.972 to 3.888 gms.)

Prep: *Extractum Eupatorii Fluidum*.

**Extractum Aconiti**—*Extract of Aconite*.

DOSE. 1-10 to  $\frac{1}{4}$  gr. (0.0064 to 0.0162 gms.)

**Extractum Aconiti Fluidum**—*Fluid Extract of Aconite*.

DOSE:  $\frac{1}{2}$  to 1 min. (0.03 to 0.06 c. c.)

**Extractum Aloes**—*Extract of Aloes*.

DOSE: 3 to 10 grs. (0.1944 to 0.648 gm.)

**Extractum Apocyni Fluidum**—*Fluid Extract of Apocynum*.

DOSE: 8 to 30 min. (0.49 to 1.85 c. c.)

**Extractum Arnice Radicis**—*Extract of Arnica Root*.

DOSE: 2 to 5 grs. (0.1296 to 0.324 gm.)

Prep: *Emplastrum Arnice*.

---

**MELLIN'S FOOD**  
Is a true Liebig's Food.

**Extractum Arnice Radicis Fluidum -***Fluid Extract of Arnica Root.*

DOSE: 5 to 20 min. (0.31 to 1.23 c. c.)

**Extractum Aromaticum Fluidum--Aromatic Fluid Extract.**

DOSE 30 to 60 min. (1.85 to 3.70 c. c.)

**Extractum Asclepiadis Fluidum--Fluid Extract of Asclepias.**

DOSE 15 to 60 min. (0.92 to 3.70 c. c.)

**Extractum Aspidospermatis Fluidum -**  
*Fluid Extract of Aspidosperma.*

DOSE. 5 to 30 min. (0.31 to 1.85 c. c.)

**Extractum Aurantii Amari Fluidum -**  
*Fluid Extract of Bitter Orange Peel.*

DOSE 15 to 60 min. (0.92 to 3.70 c. c.)

**Extractum Belladonnæ Foliorum Alcoholicum--Alcoholic Ext. Belladonna Leaves.**DOSE:  $\frac{1}{4}$  to  $\frac{3}{4}$  gr. (0.0081 to 0.0486 gm.)

Preps. Emplastrum Belladonnæ. Unguentum Belladonnæ.

**Extractum Belladonnæ Radicis Fluidum**  
*--Fluid Extract of Belladonna Root.*

DOSE: 1 to 3 min. (0.06 to 0.18 c. c.)

Prep: Linimentum Belladonnæ.

**Extractum Buchu Fluidum--Fluid Extract of Buchu.**DOSE:  $\frac{1}{4}$  to 1 fl. dr. (0.92 to 3.70 c. c.)**Extractum Calami Fluidum--Fluid Extract of Calamus.**

DOSE 15 to 60 min (0.92 to 3.70 c. c.)

**Extractum Calumbæ Fluidum--Fluid Extract of Calumba.**

DOSE: 5 to 30 min. (0.31 to 1.85 c. c.)

**Extractum Cannabis Indicæ--Extract of Indian Cannabis.**DOSE:  $\frac{1}{4}$  to 1 gr. (0.0081 to 0.0648 gm.)**Extractum Cannabis Indicæ Fluidum**  
*Fluid Extract of Indian Cannabis.*

DOSE. 2 to 5 min. (0.12 to 0.31 c. c.)

**Extractum Capsici Fluidum--Fluid Extract of Capsicum.**DOSE.  $\frac{1}{2}$  to 2 min. (0.03 to 0.12 c. c.)**Extractum Castaneæ Fluidum--Fluid Extract of Castanea.**DOSE:  $\frac{1}{2}$  to 2 fl. drs. (1.85 to 7.39 c. c.)**MELLIN'S FOOD**

Is most valuable as a nutritious food.

- Extractum Chimaphilæ Fluidum**—*Fluid Extract of Chimaphila.*  
DOSE:  $\frac{1}{2}$  to 1 fl. dr. (1.85 to 3.70 c. c.)
- Extractum Chiratæ Fluidum**—*Fluid Extract of Chirata.*  
DOSE:  $\frac{1}{2}$  to 1 fl. dr. (1.85 to 3.70 c. c.)
- Extractum Cimicifugæ**—*Extract of Cimicifuga.*  
DOSE: 3 to 5 grs. (0.1944 to 0.324 gm.)
- Extractum Cimicifugæ Fluidum**—*Fluid Extract of Cimicifuga.*  
DOSE:  $\frac{1}{2}$  to 1 fl. dr. (1.85 to 3.70 c. c.)
- Extractum Cinchonæ**—*Extract of Cinchona.*  
DOSE: 10 to 30 grs. (0.648 to 1.944 gm.)
- Extractum Cinchonæ Fluidum**—*Fluid Extract of Cinchona.*  
DOSE:  $\frac{1}{2}$  to 2 fl. drs. (1.85 to 7.39 c. c.)
- Extractum Cocæ Fluidum**—*Fluid Extract of Coca.*  
DOSE: 1 to 4 fl. drs. (3.70 to 14.79 c. c.)
- Extractum Colchici Radicis**—*Extract of Colchicum Root.*  
DOSE:  $\frac{1}{4}$  to 3 grs. (0.0162 to 0.1944 gm.)
- Extractum Colchici Radicis Fluidum**—*Fluid Extract of Colchicum Root.*  
DOSE: 2 to 5 min. (0.12 to 0.31 c. c.)
- Extractum Colchici Seminis Fluidum**—*Fluid Extract of Colchicum Seed.*  
DOSE: 1 to 5 min. (0.06 to 0.31 c. c.)
- Extractum Colocynthis**—*Extract of Colocynth.*  
DOSE:  $\frac{1}{2}$  to 2 grs. (0.13.4 to 0.1296 gm.)  
Prep: Extractum Colocynthis Compositum.
- Extractum Colocynthis Compositum**—*Compound Extract of Colocynth.*  
DOSE: 5 to 25 grs. (0.324 to 1.620 gm.)  
Preps: Pilulæ Catharticæ Compositæ. Pilulæ Catharticæ Vegetabiles.
- Extractum Conii**—*Extract of Conium.*  
DOSE:  $\frac{1}{4}$  to 2 grs. (0.0162 to 0.1296 gm.)
- Extractum Conii Fluidum**—*Fluid Extract of Conium.*  
DOSE: 1 to 10 min. (0.06 to 0.62 c. c.)
- Extractum Convallariæ Fluidum**—*Fluid Extract of Convallaria.*  
DOSE: 3 to 10 min. (0.18 to 0.62 c. c.)

---

**TRY MELLIN'S FOOD**  
For Hand-fed Infants.

**Extractum Cubebæ Fluidum—Fluid Extract of Cubeb.**

DOSE: 10 to 30 min. (0.62 to 1.85 c. c.)

**Extractum Cusso Fluidum—Fluid Extract of Cusso.**

DOSE. 1 to 4 fl. drs. (3.70 to 14.79 c. c.)

**Extractum Cypripedii Fluidum—Fluid Extract of Cypripedium.**

DOSE: 10 to 30 min. (0.62 to 1.85 c. c.)

**Extractum Digitalis—Extract of Digitalis.**DOSE:  $\frac{1}{4}$  to 1 gr. (0.0081 to 0.0648 gm.)**Extractum Digitalis Fluidum—Fluid Extract of Digitalis.**

DOSE: 1 to 2 min. (0.06 to 0.12 c. c.)

**Extractum Dulcamaræ Fluidum—Fluid Extract of Dulcamara.**DOSE  $\frac{3}{4}$  to 2 fl. drs. (2.46 to 7.39 c. c.)**Extractum Ergotæ—Extract of Ergot.**

DOSE 3 to 15 grs. (0.1944 to 0.9720 gm.)

**Extractum Ergotæ Fluidum—Fluid Extract of Ergot.**DOSE:  $\frac{1}{2}$  to 4 fl. drs. (1.85 to 4.79 c. c.)

Prep: Extractum Ergotæ.

**Extractum Eriodictyi Fluidum—Fluid Extract of Eriodictyon.**DOSE:  $\frac{1}{4}$  to 1 fl. dr. (0.92 to 3.70 c. c.)**Extractum Eucalypti Fluidum—Fluid Extract of Eucalyptus.**

DOSE. 10 to 30 min. (0.62 to 1.85 c. c.)

**Extractum Euonymi—Extract of Euonymus.**

DOSE 20 to 60 min. (1.23 to 3.70 c. c.)

**Extractum Eupatorii Fluidum—Fluid Extract of Eupatorium.**

DOSE. 15 to 60 min. (0.92 to 3.70 c. c.)

**Extractum Frangulæ Fluidum—Fluid Extract of Frangula.**DOSE  $\frac{1}{2}$  to 2 fl. drs. (1.85 to 7.39 c. c.)**Extractum Gelsemii Fluidum—Fluid Extract of Gelsemium.**

DOSE. 2 to 3 min. (0.12 to 0.18 c. c.)

**Extractum Gentianæ—Extract of Gentian.**

DOSE 5 to 10 grs. (0.324 to 0.648 gm.)

**Extractum Gentianæ Fluidum—Fluid Extract of Gentian.**

DOSE 10 to 30 min. (0.62 to 1.85 c. c.)

---

**TRY MELLIN'S FOOD**  
**For Invalids.**

- Extractum Geranii Fluidum**—*Fluid Extract of Geranium.*  
DOSE: 30 to 60 min. (1.85 to 3.70 c. c.)
- Extractum Glycyrrhizæ**—*Extract of Glycyrrhiza.*—Extract of Licorice.  
The commercial extract of the root of *Glycyrrhiza glabra*.  
DOSE: 15 to 60 grs. (0.972 to 3.888 gms.)  
Preps Trochisci Ammonii Chloridi. Trochisci Glycyrrhizæ et Opii.
- Extractum Glycyrrhizæ Fluidum**—*Fluid Extract of Glycyrrhiza.*  
DOSE:  $\frac{1}{2}$  to 1 fl. dr. (1.85 to 3.70 c. c.)
- Extractum Glycyrrhizæ Purum**—*Pure Extract of Glycyrrhiza.*  
DOSE:  $\frac{1}{4}$  to 1 dr. (0.972 to 3.888 gms.)  
Prep. Mistura Glycyrrhizæ Composita.
- Extractum Gossypii Radicis Fluidum**—*Fluid Extract of Cotton Root Bark.*  
DOSE:  $\frac{1}{2}$  to 1 fl. dr. (1.85 to 3.70 c. c.)
- Extractum Grindeliæ Fluidum**—*Fluid Extract of Grindelia.*  
DOSE:  $\frac{1}{2}$  to 1 fl. dr. (1.85 to 3.70 c. c.)
- Extractum Guaranæ Fluidum**—*Fluid Extract of Guarana.*  
DOSE:  $\frac{1}{4}$  to 1 fl. dr. (0.92 to 3.70 c. c.)
- Extractum Hæmatoxyli**—*Extract of Hæmatoxylon.*  
DOSE: 5 to 30 grs. (0.324 to 1.944 gm.)
- Extractum Hamamelidis Fluidum**—*Fluid Extract of Hamamelis.*  
DOSE:  $\frac{1}{4}$  to 1 fl. dr. (0.92 to 3.70 c. c.)
- Extractum Hydrastis Fluidum**—*Fluid Extract of Hydrastis.*  
DOSE:  $\frac{1}{2}$  to 2 fl. drs. (1.85 to 7.39 c. c.)
- Extractum Hyoscyami**—*Extract of Hyoscyamus.*  
DOSE: 1 to 3 grs. (0.0648 to 0.1944 gm.)
- Extractum Hyoscyami Fluidum**—*Fluid Extract of Hyoscyamus.*  
DOSE: 3 to 10 min. (0.18 to 0.62 c. c.)
- Extractum Ipecacuanhæ Fluidum**—*Fluid Extract of Ipecac.*  
DOSE: 3 to 8 min.,  $\frac{1}{4}$  to 1 fl. dr. (0.18 to 0.49; 0.92 to 3.70 c. c.)  
Preps Syrupus Ipecacuanhæ. Tinctura Ipecacuanhæ et Opii. Vinum Ipecacuanhæ.

**TRY MELLIN'S FOOD**  
**In Typhoid Fever.**



**Extractum Iridis—*Extract of Iris.***

DOSE 1 to 3 grs. (0.0648 to 0.1944 gm.)

**Extractum Iridis Fluidum—*Fluid Extract of Iris.***

DOSE 5 to 20 min. (0.31 to 1.23 c. c.)

**Extractum Jalapæ.**

DOSE 2 to 5 grs. (0.1296 to 0.324 gm.)

Preps Pilulæ Catharticæ Compositæ. Pilulæ Catharticæ Vegetabiles.

**Extractum Juglandis—*Extract of Juglans.***

DOSE 5 to 30 grs. (0.324 to 1.944 gm.)

**Extractum Krameriæ—*Extract of Krameria.***

DOSE 5 to 20 grs. (0.324 to 1.231 gm.)

Prep. Trochisci Krameriæ.

**Extractum Krameriæ Fluidum—*Fluid Extract of Krameria.***DOSE  $\frac{1}{4}$  to 1 fl. dr. (0.92 to 3.70 c. c.)

Prep Syrupus Krameriæ.

**Extractum Lappæ Fluidum—*Fluid Extract of Lappa.***DOSE  $\frac{1}{4}$  to 1 fl. dr. (0.92 to 3.70 c. c.)**Extractum Leptandræ—*Extract of Leptandra.***

DOSE 1 to 5 grs. (0.0648 to 0.324 gm.)

Prep. Pilulæ Catharticæ Vegetabiles.

**Extractum Leptandræ Fluidum—*Fluid Extract of Leptandra.***DOSE  $\frac{1}{4}$  to 1 fl. dr. (0.92 to 3.70 c. c.)**Extractum Lobeliæ Fluidum—*Fluid Extract of Lobelia.***

DOSE 1 to 5 min., 10 to 30 min. (0.06 to 0.31 c. c., 0.62 to 1.85 c. c.)

**Extractum Lupulini Fluidum—*Fluid Extract of Lupulin.***

DOSE 10 to 30 min. (0.62 to 1.85 c. c.)

**Extractum Matico Fluidum—*Fluid Extract of Matico.***DOSE  $\frac{1}{2}$  to 2 fl. drs. (1.85 to 7.39 c. c.)**Extractum Menispermæ Fluidum—*Fluid Extract of Menispermum.***

DOSE 5 to 30 min. (0.31 to 1.85 c. c.)

**Extractum Mezerei Fluidum—*Fluid Extract of Mezereum.***

DOSE 5 to 15 min. (0.31 to 0.92 c. c.)

Prep. Linimentum Sinapis Compositum.

**TRY MELLIN'S FOOD****In Cholera Infantum.**

**Extractum Nucis Vomicae**—*Extract of Nux Vomica.*

DOSE:  $\frac{1}{4}$  to 1 gr. (0.0162 to 0.0648 gm.)

Prep: Tinctura Nucis Vomicae.

**Extractum Nucis Vomicae Fluidum**—*Fluid Extract of Nux Vomica.*

DOSE: 1 to 4 min. (0.06 to 0.25 c. c.)

**Extractum Opii**—*Extract of Opium.*

DOSE:  $\frac{1}{4}$  to 1 gr. (0.0162 to 0.0648 gm.)

**Extractum Pareiræ Fluidum**—*Fluid Extract of Pareira.*

DOSE:  $\frac{1}{2}$  to 2 fl. drs. (1.85 to 7.39 c. c.)

**Extractum Physostigmatis** — *Extract of Physostigma.*

DOSE: 1 to  $\frac{1}{2}$  gr. (0.0064 to 0.0324 gm.)

**Extractum Phytolaccae Radicis Fluidum**—*Fluid Extract of Phytolacca Root.*

DOSE: 5 to 30 min. (0.31 to 1.85 c. c.)

**Extractum Pilocarpi Fluidum**—*Fluid Extract of Pilocarpus.*

DOSE: 10 to 30 min. (0.62 to 1.85 c. c.)

**Extractum Podophylli**—*Extract of Podophyllum.*

DOSE: 1 to 5 grs. (0.0648 to 0.324 gm.)

**Extractum Podophylli Fluidum**—*Fluid Extract of Podophyllum.*

DOSE: 5 to 15 min. (0.31 to 0.92 c. c.)

**Extractum Pruni Virginianæ Fluidum**—*Fluid Extract of Wild Cherry.*

DOSE:  $\frac{1}{2}$  to 1 fl. dr. (1.85 to 3.70 c. c.)

**Extractum Quassiae**—*Extract of Quassia.*

DOSE: 1 to 3 grs. (0.0648 to 0.1944 gm.)

**Extractum Quassiae Fluidum**—*Fluid Extract of Quassia.*

DOSE: 5 to 30 min. (0.31 to 1.85 c. c.)

**Extractum Rhamni Purshianæ Fluidum**—*Fluid Extract of Rhamnus Purshiana.*

DOSE:  $\frac{1}{4}$  to 1 fl. dr. (0.92 to 3.70 c. c.)

**Extractum Rhei**—*Extract of Rhubarb.*

DOSE: 5 to 15 grs. (0.324 to 0.972 gm.)

**Extractum Rhei Fluidum**—*Fluid Extract of Rhubarb.*

DOSE:  $\frac{1}{4}$  to 1 fl. dr. (0.92 to 3.70 c. c.)

Preps: Mistura Rhei et Sodæ. Syrupus Rhei.)

---

**TRY MELLIN'S FOOD**  
In Intestinal Indigestion.

**Extractum Rhois Glabræ Fluidum—Fluid**  
*Extract of Rhus Glabra.*DOSE  $\frac{1}{2}$  to 2 fl. drs. (1.85 to 7.39 c. c.)**Extractum Rosæ Fluidum—Fl. Ext. of Rose.**DOSE:  $\frac{1}{2}$  to 2 fl. drs. (1.85 to 7.39 c. c.)

Prep: Mel Rosæ. Syrupus Rosæ.

**Extractum Rubi Fluidum—Fluid Extract of**  
*Rubus.*DOSE  $\frac{1}{2}$  to 2 fl. drs. (1.85 to 7.39 c. c.)

Prep: Syrupus Rubi.

**Extractum Rumicis Fluidum—Fluid Ex-**  
*tract of Rumex.*DOSE  $\frac{1}{4}$  to 1 fl. dr. (0.92 to 3.70 c. c.)**Extractum Sabinæ Fluidum—Fluid Extract**  
*of Savine.*

DOSE: 5 to 15 min. (0.31 to 0.92 c. c.)

Prep: Ceratum Sabinæ.

**Extractum Sanguinariæ Fluidum—Fluid**  
*Extract of Sanguinaria.*

DOSE: 1 to 5 min. (0.06 to 0.31 c. c.)

**Extractum Sarsaparillæ Fluidum—Fluid**  
*Extract of Sarsaparilla.*DOSE:  $\frac{1}{4}$  to 1 fl. dr. (0.92 to 3.70 c. c.)

Prep: Syrupus Sarsaparillæ Compositus.

**Extractum Sarsaparillæ Fluidum Com-**  
*positum—Compound Fluid Extract of Sar-*  
*saparilla.*DOSE.  $\frac{1}{2}$  to 1½ fl. drs. (1.85 to 5.54 c. c.)**Extractum Scillæ Fluidum—Fluid Extract**  
*of Squill.*

DOSE. 1 to 5 min. (0.06 to 0.31 c. c.)

Prep: Syrupus Scillæ Compositus.

**Extractum Scoparii Fluidum—Fluid Ex-**  
*tract of Scoparius.*DOSE.  $\frac{1}{4}$  to 1 fl. dr. (0.92 to 3.70 c. c.)**Extractum Scutellarie Fluidum—Fluid**  
*Extract of Scutellaria.*DOSE:  $\frac{1}{2}$  to 2 fl. drs. (1.85 to 7.39 c. c.)**Extractum Senegæ Fluidum—Fluid Extract**  
*of Senega.*

DOSE. 3 to 15 min. (0.18 to 0.92 c. c.)

Preps Syrupus Scillæ Compositus. Syrupus Senegæ.

**Extractum Sennæ Fluidum—Fluid Extract**  
*of Senna.*

DOSE. 1 to 3 fl. drs. (3.70 to 11.09 c. c.)

Prep: Syrupus Sarsaparillæ Compositus.

---

**TRY MELLIN'S FOOD**  
**As a diet for Nursing Mothers.**

**Extractum Serpentariæ Fluidum**—*Fluid Extract of Serpentaria.*

DOSE. 1 to 4 fl. drs. (3.70 to 14.79 c. c.)

**Extractum Spigeliæ Fluidum**—*Fluid Extract of Spigelia.*

DOSE:  $\frac{1}{2}$  to 2 fl. drs. (1.85 to 7.39 c. c.)

**Extractum Stillingiæ Fluidum**—*Fluid Extract of Stillingia.*

DOSE:  $\frac{1}{4}$  to 1 fl. dr. (0.92 to 3.70 c. c.)

**Extractum Stramonii Seminis**—*Extract of Stramonium Seed*—**Extractum Stramonii**, U. S. P., 1880.

DOSE.  $\frac{1}{4}$  to  $\frac{1}{2}$  gr. (0.081 to 0.0324 gm.)

Prep: Unguentum Stramonii.

**Extractum Stramonii Seminis Fluidum**—*Fluid Extract of Stramonium Seed*—**Extractum Stramonii Fluidum**, U. S. P., 1880.

DOSE: 1 to 3 min. (0.06 to 0.18 c. c.)

**Extractum Taraxaci**—*Extract of Taraxacum.*

DOSE: 6 to 30 grs. 0.324 to 1.944 gm.)

**Extractum Taraxaci Fluidum**—*Fluid Extract of Taraxacum.*

DOSE. 1 to 3 fl. drs. (3.70 to 11.09 c. c.)

**Extractum Triticæ Fluidum**—*Fluid Extract of Triticum.*

DOSE. 1 to 4 fl. drs. (3.70 to 14.79 c. c.)

**Extractum Uvæ Ursi**—*Extract of Uva Ursi.*

DOSE: 5 to 15 grs. (0.324 to 0.972 gm.)

**Extractum Uvæ Ursi Fluidum**—*Fluid Extract of Uva Ursi.*

DOSE:  $\frac{1}{2}$  to 1 fl. dr. (1.85 to 3.70 c. c.)

**Extractum Valerianæ Fluidum**—*Fluid Extract of Valerian.*

DOSE  $\frac{1}{2}$  to 1 fl. dr. (1.85 to 3.70 c. c.)

**Extractum Veratri Viridis Fluidum**—*Fluid Extract of Veratrum Viride.*

DOSE: 1 to 5 min. (0.06 to 0.31 c. c.)

**Extractum Viburni Opuli Fluidum**—*Fluid Extract of Viburnum Opulus.*

DOSE: 1 to 2 fl. drs. (3.70 to 7.39 c. c.)

**Extractum Viburni Prunifolii Fluidum**—*Fluid Extract of Viburnum Prunifolium.*

DOSE:  $\frac{1}{4}$  to 1 fl. dr. (0.92 to 3.70 c. c.)

**Extractum Xanthoxyli Fluidum**—*Fluid Extract of Xanthoxylum.*

DOSE:  $\frac{1}{2}$  to 1 fl. dr. (1.85 to 3.70 c. c.)

## MELLIN'S FOOD

Makes cow's milk easily digestible.

**Extractum Zingiberis Fluidum**—*Fluid Extract of Ginger.*

DOSE: 10 to 30 min. (0.62 to 1.85 c. c.)

Prep: Syrupus Zingiberis.

**Fel Bovis—Oxgall—Fel Tauri.**

The fresh bile of *Bos Taurus*.

Prep Fel Bovis Purificatum.

**Fel Bovis Purificatum**—*Purified Oxgall.*

DOSE. 3 to 10 grs. (0.194 to 0.648 gm.)

**Ferri Carbonas Saccharatus**—*Saccharated Ferrous Carbonate.*

DOSE. 2 to 10 grs. (0.1296 to 0.648 gm.)

**Ferri Chloridum**—*Ferric Chloride.*

DOSE. 1 to 3 grs. (0.0648 to 0.1944 gm.)

**Ferri Citras**—*Ferric Citrate.*

DOSE 5 to 15 grs. (0.324 to 0.972 gm.)

Preps Ferri et Quininae Citras. Ferri et Quininae Citras Solubilis.

**Ferri et Ammonii Citras**—*Iron and Ammonium Citrate.*

DOSE: 5 to 15 grs. (0.324 to 0.972 gm.)

Preps Ferri et Strychninae Citras. Vinum Ferri Citratis.

**Ferri et Ammonii Sulphas**—*Iron and Ammonium Sulphate—Ammonio Ferric Sulphate. Ammonio Ferric Alum.*

DOSE: 5 to 15 grs. (0.324 to 0.972 gm.)

**Ferri et Ammonii Tartras**—*Iron and Ammonium Tartrate—Ammonio-Ferric Tartrate.*

DOSE: 10 to 30 grs. (0.648 to 1.944 gm.)

**Ferri et Potassii Tartras**—*Iron and Potassium Tartrate—Potassio-Ferric Tartrate.*

DOSE. 10 to 30 grs. (0.648 to 1.944 gm.)

**Ferri et Quininae Citras**—*Iron and Quinine Citrate.*

DOSE: 3 to 10 grs. (0.1944 to 0.648 gm.)

**Ferri et Quininae Citrate Solubilis**—*Soluble Iron and Quinine Citrate.*

DOSE. 3 to 10 grs. (0.1944 to 0.648 gm.)

Prep. Vinum Ferri Amarum.

**Ferri et Strychninae Citras**—*Iron and Strychnine Citrate.*

DOSE. 1 to 5 grs. (0.0648 to 0.324 gm.)

**Ferri Hypophosphis**—*Ferric Hypophosphite.*

DOSE 5 to 10 grs. (0.324 to 0.648 gm.)

---

**MELLIN'S FOOD**

Is readily assimilated.

**Ferri Iodidum Saccharatum—*Saccharated Ferrous Iodide.***

DOSE: 5 to 15 grs. (0.324 to 0.972 gm.)

**Ferri Lactas—*Ferrous Lactate.***

DOSE: 1 to 5 grs. (0.0648 to 0.324 gm.)

Prep. Syrupus Hypophosphitum cum Ferro.

**Ferri Oxidum Hydratum—*Ferric Hydrate—Ferric Hydroxide. Hydrated Oxide of Iron.***

DOSE: 1 to 4 drs. (3.888 to 15.552 gms.)

Preps: Emplastrum Ferri. Trochisci Ferri.

**Ferri Oxidum Hydratum cum Magnesia—*Ferric Hydrate with Magnesia.—Arsenic Antidote.***

DOSE: 1 to 4 drs. (3.888 to 15.552 gms.)

**Ferri Phosphas Solubilis—*Soluble Ferric Phosphate.***

DOSE: 5 to 10 grs. (0.324 to 0.648 gm.)

Prep. Syrupus Ferri. Quininae et Strychninae Phosphatum.

**Ferri Pyrophosphas Solubilis—*Soluble Ferric Pyrophosphate.***

DOSE: 2 to 5 grs. (0.1296 to 0.324 gm.)

**Ferri Sulphas—*Ferrous Sulphate.***

DOSE: 1 to 5 grs. (0.0648 to 0.324 gm.)

Preps: Ferri Sulphas Exsiccatus. Ferri Sulphas Granulatus.

**Ferri Sulphas Exsiccatus—*Dried Ferrous Sulphate.***DOSE:  $\frac{1}{2}$  to 3 grs. (0.324 to 0.1944 gms.)

Prep: Pilulae Aloes et Ferri.

**Ferri Sulphas Granulatus—*Granulated Ferrous Sulphate—Ferri Sulphas Præcipitatus.***

DOSE: 1 to 5 grs. (0.0648 to 0.324 gm.)

**Ferri Valerianas—*Ferric Valerianate.***DOSE:  $\frac{1}{2}$  to 2 grs. (0.0324 to 0.1296 gm.)**Ferrum—*Iron.*****Ferrum Reductum—*Reduced Iron.***

DOSE: 1 to 5 grs. (0.0648 to 0.324 gm.)

**Ficus—*Fig.***

The fleshy receptacle of Ficus Carica, bearing fruit upon its inner surface.

DOSE: Freely.

Prep: Confectio Sennæ.

---

**MELLIN'S FOOD**

Is a soluble dry extract from wheat and malt.

**Fœniculum—Fennel.**

The fruit of *Fœniculum capillaceum*.

DOSE: 15 to 30 grs. (0.972 to 1.944 gm.)

**Frangula—Frangula—Buckthorn.**

The bark of *Rhamnus Frangula*, collected at least one year before being used.

DOSE:  $\frac{1}{2}$  to 2 drs (1.944 to 7.776 gms.)

Prep: *Extractum Frangulæ Fluidum*.

**Galla—Nutmall.**

An excrescence on *Quercus lusitanica*, caused by the punctures and deposited ova of *Cynips Gallæ tinctoriæ*.

DOSE: 1 to 2 drs. (3.888 to 7.776 gms.)

Preps: *Tinctura Gallæ*. *Unguentum Gallæ*.

**Gelsemium—Gelsemium—Yellow Jasmine.**

The rhizome and roots of *Gelsemium sempervirens*.

DOSE: 5 to 10 grs. (0.324 to 0.648 gm.)

Preps: *Extractum Gelsemii Fluidum*. *Tinctura Gelsemii*.

**Gentiana—Gentian.**

The root of *Gentiana lutea*.

DOSE: 5 to 30 grs. (0.324 to 1.944 gm.)

Preps: *Extractum Gentianæ*. *Extractum Gentianæ Fluidum*. *Tinctura Gentianæ Composita*.

**Geranium—Geranium—Cranesbill.**

The rhizome of *Geranium maculatum*.

DOSE:  $\frac{1}{4}$  to 1 dr. (0.972 to 3.888 gms.)

Prep: *Extractum Geranii Fluidum*.

**Glycerinum—Glycerin.**

DOSE: 5 to 60 min. (0.31 to 3.70 c. c.)

Prep: *Suppositoria Glycerini*.

**Glyceritum Acidi Carbolicæ—Glycerite of Carbolic Acid.**

Carbolic Acid, 20%.

DOSE: 2 to 5 min. (0.12 to 0.31 c. c.)

**Glyceritum Acidi Tannici—Glycerite of Tannic Acid.**

Tannic Acid, 20%.

**Glyceritum Amyli—Glycerite of Starch.**

DOSE: Freely.

**Glyceritum Boroglycerini—Glycerite of Boroglycerin—Glycerite of Glyceryl Borate. Solution of Boroglyceride.****Glyceritum Hydrastis—Glycerite of Hydrastis.****Glyceritum Vitelli—Glycerite of Yolk of Egg—Glyconin.**

DOSE: Freely.

**MELLIN'S FOOD**

Is entirely free from starch and cane sugar.

**Glycyrrhiza—*Glycyrrhiza*—Licorice Root.**

The root of *Glycyrrhiza glabra* and of the variety *glandulifera*.

DOSE: Freely.

Preps: Extractum Glycyrrhizæ Fluidum. Extractum Glycyrrhizæ Purum. Glycyrrhizinum Ammoniatum. Pulvis Glycyrrhizæ Compositus.

**Glycyrrhizinum Ammoniatum—*Ammoniated Glycyrrhizin*.**

DOSE: 5 to 15 grs. (0.324 to 0.972 gm.)

**Gossypii Radicis Cortex—*Cotton Root Bark*.**

The bark of the root of *Gossypium herbaceum* and of other species of *Gossypium*.

DOSE:  $\frac{1}{4}$  to 1 dr. (0.972 to 3.888 gms.)

Prep: Extractum Gossypii Radicis Fluidum.

**Gossypium Purificatum—*Purified Cotton—Absorbent Cotton*.**

The hairs of the seed of *Gossypium herbaceum* and of other species of *Gossypium*, freed from adhering impurities, and deprived of fatty matter.

Prep: Pyroxylinum.

**Granatum—*Pomegranate*.**

The bark of the stem and root of *Punica Granatum*.

DOSE:  $\frac{1}{2}$  to 1½ drs. (1.944 to 5.832 gms.)

**Grindella—*Grindelia*.**

The leaves and flowering tops of *Grindelia robusta* and of *Grindelia squarrosa*.

DOSE:  $\frac{1}{4}$  to 1 dr. (0.972 to 3.888 gms.)

Prep: Extractum Grindeliæ Fluidum.

**Guaiaci Lignum—*Guaiacum Wood*.**

The heart wood of *Guaiacum officinale* Linne, and of *Guaiacum sanctum*.

DOSE:  $\frac{1}{4}$  to 1 dr. (0.972 to 3.888 gms.)

**Guaiaci Resina—*Guaiac*.**

The resin of the wood of *Guaiacum officinale*.

DOSE: 5 to 30 grs. (0.324 to 1.944 gm.)

Prep: Pilulæ Antimonii Compositæ. Tinctura Guaiaci. Tinctura Guaiaci Ammoniata.

**Guarana—*Guarana*.**

A dried paste, consisting chiefly of the crushed or pounded seeds of *Paullinia Cupana*.

DOSE:  $\frac{1}{4}$  to 1 dr. (0.972 to 3.888 gms.)

Preps: Extractum Guaranæ Fluidum.

**Hæmatoxylon—*Hæmatoxylon*—Logwood.**

The heart-wood of *Hæmatoxylon campechianum*.

Prep: Extractum Hæmatoxvli.

**MELLIN'S FOOD**

Stimulates the bowels to regular action.



**Hamamelis—Hamamelis—Witchhazel.**

The leaves of *Hamamelis virginiana*.

Prep: Extractum Hamamelidis Fluidum.

**Hedeoma—Hedeoma—Pennyroyal.**

The leaves and tops of *Hedeoma pulegioides*.

**Humulus—Hops.**

The strobiles of *Humulus Lupulus*.

Prep: Tinctura Humuli.

**Hydrargyri Chloridum Corrosivum—Corrosive Mercuric Chloride—Corrosive Chloride of Mercury. Corrosive Sublimate.**

DOSE: 1-32 to  $\frac{1}{4}$  gr. (0.0020 to 0.0081 gm.)

**Hydrargyri Chloridum Mite—Mild Mercurous Chloride—Calomel. Mild Chloride of Mercury.**

DOSE: 1 to 15 grs. (0.0648 to 0.972 gm.)

Preps: Pilule Antimoni Compositæ, Pilule Cathartice Compositæ.

**Hydrargyri Cyanidum—Mercuric Cyanide.**

DOSE: 1-24 to  $\frac{1}{2}$  gr. (0.0027 to 0.0081 gm.)

**Hydrargyri Iodidum Flavum—Yellow Mercurous Iodide—Hydrargyri Iodidum Viride, U.S.P., 1880. Protoiodide of Mercury. Yellow (or Green) Iodide of Mercury.**

DOSE:  $\frac{1}{2}$  to 3 grs. (0.0324 to 0.1944 gm.)

**Hydrargyri Iodidum Rubrum—Red Mercuric Iodide—Bin-Iodide of Mercury. Red Iodide of Mercury.**

DOSE: 1-16 to  $\frac{1}{4}$  gr. (0.0040 to 0.0216 gm.)

Prep: Liquor Arseni et Hydrargyri Iodidi.

**Hydrargyri Oxidum Flavum—Yellow Mercuric Oxide.**

Prep: Unguentum Hydrargyri Oxidi Flavi.

**Hydrargyri Oxidum Rubrum—Red Mercuric Oxide—Red Precipitate.**

Prep: Unguentum Hydrargyri Oxidi Rubri.

**Hydrargyri Subsulphas Flavus—Yellow Mercuric Subsulphate—Basic Mercuric Sulphate. Turpeth Mineral.**

DOSE: 2 to 4 grs. (0.1296 to 0.2592 gm.)

**Hydrargyrum—Mercury—Quicksilver.**

Preps: Emplastrum Ammoniac cum Hydrargyro, Emplastrum Hydrargyri, Hydrargyrum cum Creta, Massæ Hydrargyri, Unguentum Hydrargyri.

**Hydrargyrum Ammoniatum—Ammoniated Mercury—White Precipitate. Mercuric Ammonium Chloride.**

Prep: Unguentum Hydrargyri Ammoniatæ.

"With children sick with Typhoid Fever, I have found MELLIN'S an excellent food."

**Hydrargyrum cum Creta—Mercury with Chalk.**

Mercury, 38%.

DOSE: 3 to 10 grs. (0.1944 to 0.648 gm.)

**Hydrastininæ Hydrochloras—Hydrastinine Hydrochlorate.**

The hydrochlorate of an artificial alkaloid derived from Hydrastine, the latter being obtained from Hydrastis.

**Hydrastis—Hydrastis—Golden Seal.**

The rhizome and roots of *Hydrastis canadensis*.

Preps: Extractum Hydrastis Fluidum. Glyceritum Hydrastis. Tinctura Hydrastis.

**Hyoscinæ Hydrobromas—Hyoscine Hydrobromate.**

The hydrobromate of an alkaloid obtained from *Hyoscyamus*.

DOSE: 1-150 to 1-100 gr. (0.0001 to 0.00064 gm.)

**Hyoscyaminæ Hydrobromas—Hyoscyamine Hydrobromate.**

The hydrobromate of an alkaloid obtained from *Hyoscyamus*.

**Hyoscyaminæ Sulphas—Hyoscyamine Sulphate.**

The neutral sulphate of an alkaloid obtained from *Hyoscyamus*.

DOSE: 1-64 to 1-32 gr. (0.0010 to 0.0020 gm.)

**Hyoscyamus—Hyoscyamus—Henbane.**

The leaves and flowering tops of *Hyoscyamus niger*, collected from plants of the second year's growth.

DOSE: 5 to 15 grs. (0.324 to 0.972 gm.)

Preps: Extractum Hyoscyami Extractum  
Hyoscyami Fluidum. Tinctura Hyoscyami.

**Ichthyocolla—Isinglass.**

The swimming bladder of *Acipenser Huso*.

Prep: Emplastrum Ichthyocollæ.

**Illicium Illicium—Star Anise.**

The fruit of *Illicium verum*.

DOSE: 5 to 30 grs. (0.324 to 1.944 gm.)

**Infusa Infusions.**

An ordinary infusion, the strength of which is not directed by the physician nor specified by the Pharmacopœia, shall contain 5% drug.

**Infusum Cinchonæ—Infusion of Cinchona.**

Cinchona, 6%.

DOSE: 1 to 2 fl. ozs. (29.57 to 59.10 c. c.)

**Infusum Digitalis—Infusion of Digitalis.**

Digitalis, 1.5%.

DOSE: 1 to 4 fl. drs. (3.70 to 14.79 c. c.)

---

**MELLIN'S FOOD**

Contains the nitrogenous and phosphatic matters essential for the growing child.

**Infusum Pruni Virginianæ—Infusion of Wild Cherry.**

Wild Cherry, 4%.

DOSE: 1 to 2 fl. ozs. (29.57 to 59.10 c. c.)

**Infusum Sennæ Compositum—Compound Infusion of Senna—Black Draught.**

DOSE: 1 to 2½ fl. ozs. (29.57 to 88.72 c. c.)

**Inula—Inula—Elecampane.**The root of *Inula Helenum*.

DOSE: ¼ to 1 dr. (0.972 to 3.888 gms.)

**Iodoformum—Iodoform.**

DOSE: ½ to 3 grs. (0.0324 to 0.1944 gm.)

Prep: Unguentum Iodoformi.

**Iodum—Iodine.**

DOSE: ¼ to 1 gr. (0.0162 to 0.0648 gm.)

Preps: Liquor Iodi Compositus. Tinctura Iodi. Unguentum Iodi.

**Ipecacuanha—Ipecac.**The root of *Cephaelis Ipecacuanha*.

DOSE: ½ to 30 grs. (0.0324 to 1.944 gm.)

Preps: Extractum Ipecacuanhæ Fluidum. Pulvis Ipecacuanhæ et Opii. Trochisci Ipecacuanhæ. Trochisci Morphine et Ipecacuanhæ.

**Iris—Iris—Blue Flag.**The rhizome and roots of *Iris versicolor*.

DOSE: 10 to 30 grs. (0.648 to 1.944 gm.)

Preps: Extractum Iridis. Extractum Iridis Fluidum.

**Jalapa—Jalap.**The tuberous root of *Ipomœa Jalapa*.

DOSE: 10 to 30 grs. (0.648 to 1.944 gm.)

Preps: Extractum Jalapæ Alcoholicum. Pulvis Jalapæ Compositus. Resina Jalapæ.

**Juglans—Juglans—Butternut.**The bark of the root of *Juglans cinerea*.

DOSE: 1 to 2 drs. (3.888 to 7.776 gms.)

Prep: Extractum Juglandis.

**Kamala—Kamala—Rottlera.**The glands and hairs from the capsules of *Mallotus philippinensis*.

DOSE: 1 to 2 drs. (3.888 to 7.776 gm.)

**Kino—Kino.**The inspissated juice of *Pterocarpus Marsupium*

DOSE: 5 to 30 grs. (0.324 to 1.944 gm.)

Prep: Tinctura Kino.

---

**MELLIN'S FOOD**Prepared with milk fulfils every requirement  
of a food for infants.

**Krameria—Krameria—Rhatany.**

The root of *Krameria triandra* and of *Krameria Ixina*.

DOSE: 5 to 30 grs. (0.324 to 1.944 gm.)

Preps: Extractum *Krameria*. Extractum *Krameria*  
Fluidum. Tinctura *Krameria*.

**Lactucarium Lactucarium.**

The concrete milk juice of *Lactuca virosa*.

DOSE: 5 to 60 grs. (0.324 to 3.888 gms.)

Prep: Tinctura *Lactucarii*.

**Lappa—Lappa—Burdock.**

The root of *Arctium Lappa*, and of other species of *Arctium*.

DOSE  $\frac{1}{2}$  to 1 dr. (1.944 to 3.888 gms.)

Prep: Extractum *Lappæ* Fluidum.

**Leptandra—Leptandra—Culver's Root.**

The rhizome and roots of *Veronica virginica*.

DOSE:  $\frac{1}{4}$  to 1 dr. (0.972 to 3.888 gms.)

Preps: Extractum *Leptandræ*. Extractum *Leptandræ*  
Fluidum.

**Limonis Cortex—Lemon Peel.**

The rind of the recent fruit of *Citrus Limonum*.

Prep: Spiritus *Limonis*.

**Limonis Succus—Lemon Juice.**

The freshly expressed juice of the ripe fruit of *Citrus Limonum*.

**Linimentum Ammonia—Ammonia Liniment—Volatile Liniment.****Linimentum Belladonnæ—Belladonna Liniment.****Linimentum Calcis—Lime Liniment. Carron Oil.****Linimentum Camphoræ—Camphor Liniment.****Linimentum Chloroformi—Chloroform Liniment.****Linimentum Saponis—Soap Liniment.**

Prep: Linimentum *Chloroformi*.

**Linimentum Saponis Mollis—Liniment of Soft Soap—Tinctura Saponis Viridis. U. S. P., 1880.****Linimentum Sinapis Compositum—Compound Liniment of Mustard.****Linimentum Terebinthinæ—Turpentine Liniment.****Linum—Linseed—Flaxseed.**

The seed of *Linum Usitatissimum*.

**MELLIN'S FOOD**

Contains a large amount of proteids and soluble phosphates.

**Liquor Acidi Arsenosi—Solution of Arsenous Acid.**

Arsenous Acid, 1%.

DOSE: 3 to 10 min. (0.18 to 0.62 c. c.)

**Liquor Ammonii Acetatis—Solution of Ammonium Acetate—Spirit of Mindererus.**

An aqueous solution of Ammonium Acetate, containing about 7% of the salt, together with small amounts of acetic and carbonic acids.

DOSE:  $\frac{1}{4}$  to 1 fl. oz. (7.39 to 29.57 c. c.)

Prep: Liquor Ferri et Ammonii Acetatis.

**Liquor Arseni et Hydrargyri Iodidi—Solution of Arsenic and Mercury Iodide. Donovan's Solution.**

Arsenic Iodide, 1%. Red Mercuric Iodide, 1%.

DOSE: 3 to 15 min. (0.18 to 0.92 c. c.)

**Liquor Calcis—Solution of Lime—Solution of Calcium Hydrate. Lime Water.**

A saturated aqueous solution of Calcium Hydrate.

DOSE:  $\frac{1}{2}$  to 4 fl. ozs. (14.79 to 118.29 c. c.)

Prep: Linimentum Calcis.

**Liquor Ferri Acetatis—Solution of Ferric Acetate.**

An aqueous solution of Ferric Acetate, containing about 31% of the anhydrous salt, and representing about 7.5% of metallic iron.

DOSE: 2 to 10 min. (0.12 to 0.62 c. c.)

**Liquor Ferri Chloridi—Solution of Ferric Chloride.**

An aqueous solution of Ferric Chloride, containing about 37.8% of the anhydrous salt, corresponding to 62.9% of the crystallized salt, or to about 13% of metallic iron.

DOSE: 3 to 10 min. (0.18 to 0.62 c. c.)

Prep: Tinctura Ferri Chloridi.

**Liquor Ferri Citratis—Solution of Ferric Citrate.**

An aqueous solution of Ferric Citrate, corresponding to about 7.5% of metallic iron.

DOSE: 5 to 15 min. (0.31 to 0.92 c. c.)

Preps: Ferri Citras. Ferri et Ammonii Citras.

**Liquor Ferri et Ammonii Acetatis—Solution of Iron and Ammonium Acetate—Mistura Ferri et Ammonii Acetatis. U. S. P., 1880. Basham's Mixture.**DOSE:  $\frac{1}{4}$  to 1 fl. oz. (7.39 to 29.57 c. c.)**MELLIN'S FOOD**

Actually assists to digest milk or other foods with which it is mixed.

**Liquor Ferri Nitratis**—*Solution of Ferric Nitrate.*

An aqueous solution of Ferric Nitrate, containing about 6.2% of the anhydrous salt, and corresponding to about 1.4% of metallic iron.

DOSE: 5 to 15 min. (0.31 to 0.92 c. c.)

**Liquor Ferri Subsulphatis**—*Solution of Ferric Subsulphate*—*Solution of Basic Ferric Sulphate.* Monsel's Solution.

An aqueous solution of Basic Ferric Sulphate (of variable chemical composition), corresponding to about 13.6% of metallic iron.

DOSE: 2 to 10 min. (0.12 to 0.62 c. c.)

**Liquor Ferri Tersulphatis**—*Solution of Ferric Sulphate.*

An aqueous solution of normal Ferric Sulphate, containing about 28.7% of the salt, and corresponding to about 8% of metallic iron.

**Liquor Hydrargyri Nitratis**—*Solution of Mercuric Nitrate.*

A liquid containing about 60% of Mercuric Nitrate, together with about 11% of free Nitric Acid.

**Liquor Iodi Compositus**—*Compound Solution of Iodine*—Lugol's Solution.

DOSE: 1 to 10 min. (0.06 to 0.62 c. c.)

**Liquor Magnesii Citratis**—*Solution of Magnesium Citrate.*

DOSE: 2 to 8 fl. oz. (59.10 to 236.58 c. c.)

**Liquor Plumbi Subacetatis**—*Solution of Lead Subacetate.*

An aqueous liquid, containing in solution about 25% of Lead Subacetate.

Preps: Ceratum Plumbi Subacetatis. Liquor Plumbi Subacetatis Dilutus.

**Liquor Plumbi Subacetatis Dilutus**—*Diluted Solution of Lead Subacetate*—*Lead Water.*

**Liquor Potassæ**—*Solution of Potassa*—*Solution of Potassium Hydrate.*

An aqueous solution of Potassium Hydrate, containing about 5% of the hydrate.

DOSE: 10 to 30 min. (0.62 to 1.85 c. c.)

**Liquor Potassii Arsenitis**—*Solution of Potassium Arsenite*—Fowler's Solution.

Arsenous Acid, 1%.

DOSE: 3 to 10 min. (0.18 to 0.62 c. c.)

---

**MELLIN'S FOOD**

Has proved its greater nutritive value over all other artificial foods.

**Liquor Potassii Citratis**—*Solution of Potassium Citrate*—*Mistura Potassii Citratis*.

An aqueous liquid, containing in solution about 9% of anhydrous Potassium Citrate, together with small amounts of citric and carbonic acids.

DOSE: 1 to 8 fl. drs. (3.70 to 29.57 c. c.)

**Liquor Sodæ**—*Solution of Soda*—*Solution of Sodium Hydrate*.

An aqueous solution of Sodium Hydrate, containing about 5% of the hydrate.

DOSE: 5 to 20 min. (0.31 to 1.23 c. c.)

**Liquor Sodæ Chloratæ**—*Solution of Chlorinated Soda*—*Labarraque's Solution*.

An aqueous solution of several chlorine compounds of sodium, containing at least 2.6% by weight of available chlorine.

DOSE:  $\frac{1}{2}$  to 1 fl. dr. (1.85 to 3.70 c. c.)

**Liquor Sodii Arsenatis**—*Solution of Sodium Arsenate*.

Sodium Arsenate, 1%.

DOSE: 3 to 10 min. (0.18 to 0.62 c. c.)

**Liquor Sodii Silicatis**—*Solution of Sodium Silicate*.**Liquor Zinci Chloridi**—*Solution of Zinc Chloride*.

An aqueous solution of Zinc Chloride, containing about 50% by weight of the salt.

**Lithii Benzoas**—*Lithium Benzoate*.

DOSE: 10 to 30 grs. (0.648 to 1.944 gm.)

**Lithii Bromidum**—*Lithium Bromide*.

DOSE: 5 to 30 grs. (0.324 to 1.944 gm.)

**Lithii Carbonas**—*Lithium Carbonate*.

DOSE: 3 to 15 grs. (0.1944 to 0.972 gm.)

**Lithii Citras**—*Lithium Citrate*.

DOSE: 10 to 30 grs. (0.648 to 1.944 gms.)

**Lithii Citras Effervescens**—*Effervescent Lithium Citrate*.**Lithii Salicylas**—*Lithium Salicylate*.

DOSE: 10 to 30 grs. (0.648 to 1.944 gm.)

**Lobelia**—*Lobelia*.

The leaves and tops of *Lobelia inflata*, collected after a portion of the capsules have been inflated.

DOSE: 1 to 10 grs. (0.0648 to 0.648 gm.)

Preps: Extractum Lobeliæ Fluidum. Tinctura Lobeliæ.

**MELLIN'S FOOD**

Is a most useful addition to the diet of healthy children.

**Lupulinum—*Lupulin*.**

The glandular powder separated from the strobiles of *Humulus Lupulus*.

DOSE: 5 to 15 grs. (0.324 to 0.972 gm.)

Preps: Extractum Lupulini Fluidum Oleoresina Lupulini.

**Lycopodium—*Lycopodium*.**

The spores of *Lycopodium clavatum* and of other species of *Lycopodium*.

**Macis—*Mace*.**

The arilode of the seed of *Myristica fragrans*.

DOSE: 5 to 20 grs. (0.324 to 1.296 gm.)

**Magnesia—*Magnesia*—Light Magnesia. Calcined Magnesia.**

DOSE: 5 to 60 grs. (0.324 to 3.888 gms.)

Prep: Pulvis Rhei Compositus.

**Magnesia Ponderosa—*Heavy Magnesia*.**

DOSE: 5 to 60 grs. (0.324 to 3.888 gms.)

**Magnesii Carbonas—*Magnesium Carbonate*.**

DOSE: 1 to 3 drs. (3.888 to 11.664 gms.)

**Magnesii Citras Effervescens—*Effervescent Magnesium Citrate*.**

DOSE:  $\frac{1}{4}$  to 1 oz. (7.776 to 31.103 gms.)

**Magnesii Sulphas—*Magnesium Sulphate*—Epsom Salt.**

DOSE:  $\frac{1}{2}$  to  $1\frac{1}{2}$  ozs. (15.552 to 46.655 gms.)

Prep: Infusum Sennæ Compositum.

**Mangani Dioxidum—*Manganese Dioxide*—Mangani Oxidum Nigrum, U. S. P., 1880. Black Oxide of Manganese.**

Native crude Manganese Dioxide, containing at least 66% of the pure dioxide.

DOSE: 8 to 15 grs. (0.1944 to 0.972 gm.)

**Mangani Sulphas—*Manganese Sulphate*—Manganous Sulphate.**

DOSE: 5 to 15 grs. (0.324 to 0.972 gm.)

**Manna—*Manna*.**

Concrete, saccharin exudation of *Fraxinus Ornus*.

DOSE:  $\frac{1}{2}$  to 2 ozs. (15.552 to 62.207 gms.)

Prep: Infusum Sennæ Compositum.

**Marrubium—*Marrubium*—Horehound.**

The leaves and tops of *Marrubium vulgare*.

DOSE: 1 to 2 drs. (3.888 to 7.776 gms.)

**Massa Copaibæ—*Mass of Copaiba*—Solidified Copaiba.**

DOSE: 5 to 30 grs. (0.324 to 1.944 gm.)

**MELLIN'S FOOD**

Is unrivalled as an artificial food for children.



**Massa Ferri Carbonatis**—*Mass of Ferrous Carbonate*—Vallet's Mass.

DOSE. 3 to 7 grs. (0.1944 to 0.4336 gm.)

**Massa Hydrargyri**—*Mass of Mercury*—*Pilula Hydrargyri*. Blue Mass. Blue Pill.

DOSE. 3 to 10 grs. (0.1944 to 0.648 gm.)

**Mastiche**—*Mastic*.

A concrete, resinous exudation from *Pistacia Lentiscus*.

Prep. *Pilula Aloes et Mastiches*.

**Matico**—*Matico*.

The leaves of *Piper angustifolium*.

DOSE:  $\frac{1}{2}$  to 2 drs. (1.944 to 7.776 gms.)

Preps. *Extractum Matico Fluidum*. *Tinctura Matico*.

**Matricaria** *Matricaria*—German Chamomile.

The flower heads of *Matricaria Chamomilla*.

DOSE:  $\frac{1}{4}$  to 1 dr. (0.972 to 3.888 gms.)

**Mel**—*Honey*.

A saccharine substance deposited in the honey comb by *Apis mellifica*.

Prep. *Mel Despumatum*.

**Mel Despumatum**—*Clarified Honey*.

**Mel Rosæ**—*Honey of Rose*.

**Melissa**—*Melissa*—Balm.

The leaves and tops of *Melissa officinalis*.

DOSE. 1 to 2 drs. (3.888 to 7.776 gms.)

**Menispermum**—*Menispermum*—Yellow Parilla.  
Canadian Moonseed.

The rhizome and roots of *Menispermum canadense*.

DOSE: 5 to 30 grs (0.324 to 1.944 gm.)

Prep: *Extractum Menispermi Fluidum*.

**Mentha Piperita** *Peppermint*.

The leaves and tops of *Mentha Piperita*.

Prep: *Spiritus Menthae Piperitæ*.

**Mentha Viridis** *Spearmint*.

The leaves and tops of *Mentha viridis*.

Prep: *Spiritus Menthae Viridis*.

**Menthol**—*Menthol*.

A stearopten (having the character of a secondary alcohol), obtained from the official oil of peppermint or from Japanese or Chinese oil of peppermint.

DOSE: 1 to 5 grs. (0.0648 to 0.324 gm.)

**Methyl Salicylas** *Methyl Salicylate*—Artificial (or Synthetic) Oil of Wintergreen.

Methyl Salicylate, produced synthetically.

DOSE: 1 to 5 min. (0.06 to 0.31 c. c.)

**MELLIN'S FOOD**  
Is a true Liebig's Food.

**Mezereum—Mezereum.**

The bark of *Daphne Mezereum* and of other species of *Daphne*.

DOSE: 5 to 15 grs.

Preps: Decoctum Sarsaparillæ Compositum. Extractum Sarsaparillæ Fluidum Compositum. Extractum Mezerei Fluidum.

**Mistura Cretæ—Chalk Mixture.**

DOSE:  $\frac{1}{4}$  to 1 fl. oz. (7.39 to 29.57 c. c.)

**Mistura Ferri Composita—Compound Iron Mixture—Griffith's Mixture.**

DOSE:  $\frac{1}{2}$  to 2 fl. ozs. (14.79 to 59.10 c. c.)

**Mistura Glycyrrhizæ Composita—Compound Mixture of Glycyrrhiza. Brown Mixture.**

DOSE: 2 to 6 fl. drs. (7.39 to 22.18 c. c.)

**Mistura Rhei et Sodæ—Mixture of Rhubarb and Soda.**

DOSE:  $\frac{1}{2}$  to 2 fl. ozs. (14.79 to 59.10 c. c.)

**Morphina—Morphine.**

An alkaloid obtained from Opium.

DOSE: 1-16 to  $\frac{1}{4}$  gr. (0.0049 to 0.0162 gm.)

**Morphinæ Acetas—Morphine Acetate.**

DOSE: 1-16 to  $\frac{1}{4}$  gr. (0.0049 to 0.0216 gm.)

**Morphinæ Hydrochloras—Morphine Hydrochlorate.**

DOSE: 1-12 to  $\frac{1}{2}$  gr. (0.0054 to 0.0324 gm.)

**Morphinæ Sulphas—Morphine Sulphate.**

DOSE: 1-12 to  $\frac{1}{2}$  gr. (0.0054 to 0.0324 gm.)

Preps: Pulvis Morphinæ Compositus. Trochisci Morphinæ et Ipecacuanæ.

**Moschus—Musk.**

The dried secretion from the preputial follicles of *Moschus moschiferus*.

DOSE: 2 to 10 grs. (0.1296 to 0.648 gm.)

Prep: Tinctura Moschi.

**Mucilago Acaciæ—Mucilage of Acacia.**

Acacia, 34%.

Prep: Syrupus Acaciæ.

**Mucilago Sassafras Medullæ—Mucilage of Sassafras Pith.****Mucilago Tragacanthæ—Mucilage of Tragacanth.**

Tragacanth, 6%.

**Mucilago Ulmi—Mucilage of Elm.**

Elm, 6%.

**MELLIN'S FOOD**

Is most valuable as a nutritious food.

**Myristica—Nutmeg.**

The seed of *Myristica fragrans* deprived of its testa.

DOSE: 5 to 15 grs. (0.324 to 0.972 gm.)

Preps: Pulvis Aromaticus. Tinctura Lavandulæ Composita.

**Myrrha—Myrrh.**

A gum resin, obtained from *Commiphora Myrrha*.

DOSE: 5 to 30 grs. (0.324 to 1.944 gm.)

Preps: Mistura Ferri Composita. Pilulæ Aloes et Myrrhæ. Tinctura Aloes et Myrrhæ. Tinctura Myrrhæ.

**Naphtalinum—Naphtalin—Naphtalene.**

A hydrocarbon obtained from cold tar.

DOSE: 2 to 10 grs. (0.1296 to 0.648 gm.)

**Naphtol—Naphtol—Beta Naphtol.**

A phenol occurring in coal tar, but usually prepared artificially from Naphtalin.

DOSE: 2 to 15 grs. (0.1296 to 0.972 gm.)

**Nux Vomica—Nux Vomica.**

The seed of *Strychnos Nux-vomica*.

DOSE: 1 to 4 grs. (0.0648 to 0.2592 gm.)

Preps: Extractum Nucis Vomicae. Extractum Nucis Vomicae Fluidum.

**Oleatum Hydrargyri—Oleate of Mercury.**

Yellow Mercury Oxide, 20%.

**Oleatum Veratrinæ—Oleate of Veratrine.**

Veratrine, 2%.

**Oleatum Zinci—Oleate of Zinc.**

Zinc Oxide, 5%.

**Oleoresina Aspidii—Oleoresin of Aspidium.**

DOSE:  $\frac{1}{2}$  to 1 fl. dr. (1.85 to 3.70 c. c.)

**Oleoresina Capsici—Oleoresin of Capsicum.**

DOSE:  $\frac{1}{4}$  to 1 min. (0.015 to 0.06 c. c.)

Prep: Emplastrum Capsici.

**Oleoresina Cubebæ—Oleoresin of Cubeb.**

DOSE: 5 to 30 min. (0.31 to 1.85 c. c.)

Prep: Trochisci Cubebæ.

**Oleoresina Lupulini—Oleoresin of Lupulin.**

DOSE: 3 to 7 min. (0.15 to 0.43 c. c.)

**Oleoresina Piperis—Oleoresin of Pepper.**

DOSE:  $\frac{1}{4}$  to 2 min. (0.015 to 0.12 c. c.)

**Oleoresina Zingiberis—Oleoresin of Ginger.**

DOSE:  $\frac{1}{2}$  to 2 min. (0.03 to 0.12 c. c.)

**Oleum Adipis—Lard Oil.**

A fixed oil expressed from lard at a low temperature.

**Oleum Æthereum—Ethereal Oil.**

A volatile liquid, consisting of equal volumes of Heavy Oil of Wine and Ether.

Prep: Spiritus Ætheris Compositus.

**TRY MELLIN'S FOOD**  
For Hand-fed Infants.

**Oleum Amygdalæ Amaræ—Oil of Bitter Almond.**

A volatile oil obtained from Bitter Almond by maceration with water and subsequent distillation.

DOSE:  $\frac{1}{4}$  to  $\frac{1}{2}$  min. (0.01 to 0.03 c. c.)

Preps: Aqua Amygdalæ Amaræ. Spiritus Amygdalæ Amaræ.

**Oleum Amygdalæ Expressum—Expressed Oil of Bitter Almond.**

A fixed oil expressed from Bitter or Sweet Almond.

DOSE: 1 to 4 fl. dr. (3.70 to 14.79 c. c.)

Prep: Unguentum Aquæ Rosæ.

**Oleum Anisi—Oil of Anise.**

A volatile oil distilled from Anise.

DOSE: 1 to 5 min. (0.06 to 0.31 c. c.)

Preps: Aqua Anisi. Spiritus Aurantii Compositus. Tinctura Opii Camphorata.

**Oleum Aurantii Corticis—Oil of Orange Peel.**

A volatile oil obtained by expression from the fresh peel of either the Bitter Orange, *Citrus Vulgaris*, or the Sweet Orange, *Citrus Aurantium*.

Preps: Spiritus Aurantii. Spiritus Aurantii Compositus. Spiritus Myrciæ.

**Oleum Aurantii Florum—Oil of Orange Flowers—Oil of Neroli.**

A volatile oil distilled from the fresh flowers of the Bitter Orange, *Citrus Vulgaris*.

**Oleum Bergamottæ—Oil of Bergamot—Oleum Bergamii, U. S. P., 1880.**

A volatile oil obtained by expression from the rind of the fresh fruit of *Citrus Bergamia*.

**Oleum Betulæ Volatile—Volatile Oil of Betula—Oil of Sweet Birch.**

A volatile oil, identical with synthetic Methyl Salicylate, and almost wholly identical with Oil of Gaultheria obtained by distillation from the bark of Sweet Birch, *Betula lenta*.

DOSE: 1 to 5 min. (0.06 to 0.31 c. c.)

**Oleum Cadinum—Oil of Cade—Oleum Juniperi Empyreumaticum.**

A product of the dry distillation of the wood *Juniperus Oxycedrus*.

**Oleum Cajuputi—Oil of Cajuput.**

A volatile oil distilled from the leaves of *Melaleuca Leucadendron*.

DOSE: 1 to 5 min. (0.06 to 0.31 c. c.)

---

**TRY MELLIN'S FOOD**  
**For Invalids.**

**Oleum Carl—Oil of Caraway.**

A volatile oil distilled from Caraway.

DOSE. 1 to 5 min. (0.06 to 0.31 c. c.)

Prep Spiritus Juniperi Compositus.

**Oleum Caryophylli—Oil of Cloves.**

A volatile oil distilled from Cloves.

DOSE. 1 to 5 min. (0.06 to 0.31 c. c.)

**Oleum Chenopodii—Oil of Chenopodium—  
Oil of American Wormseed.**

A volatile oil distilled from Chenopodium.

DOSE: 3 to 10 min. (0.18 to 0.62 c. c.)

**Oleum Cinnamomi—Oil of Cinnamon—Oil of  
Cassia.**

A volatile oil distilled from Cassia Cinnamon.

DOSE. 1 to 5 min. (0.06 to 0.31 c. c.)

Preps Aqua Cinnamomi. Spiritus Cinnamomi.

**Oleum Copaibæ—Oil of Copaiba.**

A volatile oil distilled from Copaiba.

DOSE. 5 to 15 min. (0.31 to 0.92 c. c.)

**Oleum Coriandri—Oil of Coriander.**

A volatile oil distilled from Coriander.

DOSE. 2 to 5 min. (0.12 to 0.31 c. c.)

Prep Spiritus Aurantii Compositus.

**Oleum Cubebæ—Oil of Cubeb.**

A volatile oil distilled from Cubeb.

DOSE: 5 to 15 min. (0.31 to 0.92 c. c.)

**Oleum Erigerontis—Oil of Erigeron—Oil of  
Fleabane.**A volatile oil distilled from the fresh flowering herb of *Erigeron canadense*.

DOSE 5 to 30 min. (0.31 to 1.85 c. c.)

**Oleum Eucalypti—Oil of Eucalyptus.**A volatile oil distilled from the fresh leaves of *Eucalyptus globulus* and some other species of *Eucalyptus*.

DOSE: 5 to 15 min. (0.31 to 0.92 c. c.)

**Oleum Feniculi—Oil of Fennel.**

A volatile oil distilled from Fennel.

DOSE: 1 to 5 min. (0.06 to 0.31 c. c.)

Preps Aqua Feniculi. Pulvis Glycyrrhizæ Compositus.  
Spiritus Juniperi Compositus.**Oleum Gaultheriæ—Oil of Wintergreen.**A volatile oil distilled from the leaves of *Gaultheria procumbens*, consisting almost entirely of Methyl Salicylate, and nearly identical with the Volatile Oil of Betula.

DOSE: 5 to 30 min. (0.31 to 1.85 c. c.)

Prep: Spiritus Gaultheriæ.

**TRY MELLIN'S FOOD**  
**In Typhoid Fever.**

**Oleum Gossypil Seminis—Cotton Seed Oil.**

A fixed oil expressed from the seed of *Gossypium herbaceum* and of other species of *Gossypium*, and subsequently purified.

**Oleum Hedeomæ—Oil of Hedeoma—Oil of Pennyroyal.**

A volatile oil distilled from *Hedeoma*.

DOSE: 1 to 5 min. (0.06 to 0.31 c. c.)

**Oleum Juniperi—Oil of Juniper.**

A volatile oil distilled from the fruit of *Juniperus communis*.

DOSE: 5 to 15 min. (0.31 to 0.92 c. c.)

Preps: *Spiritus Juniperi*. *Spiritus Juniperi Compositus*.

**Oleum Lavandulæ Florum—Oil of Lavender Flowers.**

A volatile oil distilled from the fresh flowers of *Lavandula officinalis*.

DOSE: 1 to 5 min. (0.06 to 0.31 c. c.)

Preps: *Spiritus Lavandulæ*. *Tinctura Lavandulæ Composita*.

**Oleum Limonis—Oil of Lemon.**

A volatile oil obtained by expression from fresh Lemon Peel.

Preps: *Spiritus Aurantii Compositus*. *Spiritus Limonis*.

**Oleum Lini—Linseed Oil—Oil of Flaxseed.**

A fixed oil expressed from Linseed without the use of heat.

DOSE:  $\frac{1}{2}$  to 8 fl. drs. (1.85 to 29.57 c. c.)

Prep: *Linimentum Calcis*.

**Oleum Menthæ Piperitæ—Oil of Peppermint.**

A volatile oil expressed from Peppermint.

DOSE: 1 to 5 min. (0.06 to 0.31 c. c.)

Preps: *Aqua Menthæ Piperitæ*. *Spiritus Menthæ Piperitæ*. *Trochisci Menthæ Piperitæ*.

**Oleum Menthæ Viridis—Oil of Spearmint.**

A volatile oil distilled from Spearmint.

DOSE: 1 to 5 min. (0.06 to 0.31 c. c.)

Preps: *Aqua Menthæ Viridis*. *Spiritus Menthæ Viridis*.

**Oleum Morrhuæ—Cod Liver Oil—Oleum Jecoris Aselli.**

A fixed oil obtained from the fresh livers of *Gadus Morrhua* and of other species of *Gadus*.

DOSE:  $\frac{1}{4}$  to 1 fl. oz. (7.39 to 29.57 c. c.)

**Oleum Myrciæ—Oil of Myrcia—Oil of Bay.**

A volatile oil distilled from the leaves of *Myrcia acris*.

Prep: *Spiritus Myrciæ*.

**Oleum Myristicæ—Oil of Nutmeg.**

A volatile oil distilled from Nutmeg.

DOSE: 1 to 3 min. (0.06 to 0.18 c. c.)

Prep: *Spiritus Myristicæ*.

---

TRY MELLIN'S FOOD  
In Cholera Infantum.

**Oleum Olivæ—Olive Oil.**

A fixed oil expressed from the ripe fruit of *Olea europæa*.

**Oleum Phosphoratum—Phosphorated Oil.**

Phosphorus, 1%.

DOSE: 1 to 5 min. (0.06 to 0.31 c. c.)

**Oleum Picis Liquidæ—Oil of Tar.**

A volatile oil distilled from tar.

**Oleum Pimentæ—Oil of Pimenta—Oil of Allspice.**

A volatile oil distilled from Pimenta.

DOSE: 1 to 5 min. (0.06 to 0.31 c. c.)

Prep: Spiritus Myrciæ.

**Oleum Ricini—Castor Oil.**

A fixed oil expressed from the seed of *Ricinus communis*.

DOSE:  $\frac{1}{4}$  to 2 fl. ozs. (7.39 to 59.10 c. c.)

**Oleum Rosæ—Oil of Rose.**

A volatile oil distilled from the fresh flowers of *Rosa damascena*.

**Oleum Rosmarini—Oil of Rosemary.**

A volatile oil distilled from the leaves of *Rosmarinus officinalis*.

DOSE: 1 to 5 min. (0.06 to 0.31 c. c.)

Prep: Tinctura Lavandulæ Composita.

**Oleum Sabinæ—Oil of Savine.**

A volatile oil distilled from Savine.

DOSE: 1 to 5 min. (0.06 to 0.31 c. c.)

**Oleum Santali—Oil of Santal—Oil of Sandalwood.**

A volatile oil distilled from the wood of *Santalum album*.

DOSE: 10 to 30 min. (0.62 to 1.85 c. c.)

**Oleum Sassafras—Oil of Sassafras.**

A volatile oil distilled from Sassafras.

DOSE: 1 to 5 min. (0.06 to 0.31 c. c.)

**Oleum Sesami—Oil of Sesamum—Sesame Oil. Teel Oil. Benne Oil.**

A fixed oil expressed from the seed of *Sesamum indicum*.

DOSE: Freely.

**Oleum Sinapis Volatile—Volatile Oil of Mustard.**

A volatile oil obtained from Black Mustard by maceration with water and subsequent distillation.

DOSE:  $\frac{1}{4}$  to  $\frac{1}{2}$  min. (0.0075 to 0.015 c. c.)

Prep: Linimentum Sinapis Compositum.

**Oleum Terebinthinæ—Oil of Turpentine.**

A volatile oil distilled from Turpentine.

DOSE: 5 min. to 2 fl. drs. (0.31 to 7.39 c. c.)

Preps: Linimentum Terebinthinæ. Oleum Terebinthinæ Rectificatum.

**TRY MELLIN'S FOOD**  
In Intestinal Indigestion.

**Oleum Terebinthinæ Rectificatum—Rectified Oil of Turpentine.**

DOSE: 5 to 30 min. (0.31 to 1.85 c. c.)

**Oleum Theobromatis—Oil of Theobroma—Oleum Theobromæ, U. S. P., 1880. Butter of Cacao.**

A fixed oil expressed from the seed of *Theobroma Cacao*.

**Oleum Thymi—Oil of Thyme.**

A volatile oil distilled from the leaves and flowering tops of *Thymus vulgaris*.

DOSE: 1 to 5 min. (0.06 to 0.31 c. c.)

**Oleum Tigllii—Croton Oil.**

A fixed oil expressed from the seed of *Croton Tiglium*.

DOSE:  $\frac{1}{2}$  to 2 min. (0.03 to 0.12 c. c.)

**Opil Pulvis—Powdered Opium.**

Opium dried at a temperature not exceeding 85 degrees C. (185 deg. F.), and reduced to a very fine (No. 80) powder.

DOSE:  $\frac{1}{4}$  to 2 grs. (0.0162 to 0.1296 gm.)

Preps: *Acetum Opil*. *Extractum Opil*. *Opium Deodoratum*. *Pilulæ Opil*. *Pulvis Ipecacuanhæ et Opil*. *Tinctura Opil*. *Tinctura Opil Camphorata*. *Tinctura Opil Deodorati*. *Trochisci Glycyrrhizæ et Opil*. *Vinum Opil*.

**Opium—Opium.**

The concrete, milky exudation obtained by incising the unripe capsules of *Papaver somniferum*, and yielding in its normal moist condition not less than 9% of crystallized morphine, when assayed by the official process.

DOSE:  $\frac{1}{4}$  to 2 grs. (0.0162 to 0.1296 gm.)

**Opium Deodoratum—Deodorized Opium—Opium Denarcotisatum, U. S. P., 1880.**

DOSE:  $\frac{1}{4}$  to 2 grs. (0.0162 to 0.1296 gm.)

**Pancreatinum—Pancreatin.**

A mixture of the enzymes naturally existing in the pancreas of warm-blooded animals, usually obtained from the fresh pancreas of the hog.

DOSE: 5 to 15 grs. (0.324 to 0.972 gm.)

**Paraldehydum—Paraldehyde.**

A polymeric form of Ethylic Aldehyde.

DOSE: 1 to 3 grs. (0.0648 to 0.1944 gm.)

**Pareira—Pareira—Pareira Brava.**

The root of *Chondodendron tomentosum*.

DOSE:  $\frac{1}{2}$  to 1 dr. (1.944 to 3.888 gms.)

Prep: *Extractum Pareiræ Fluidum*.

**Pepo—Pumpkin Seed.**

The seed of *Cucurbita Pepo*.

DOSE: 1 to 3 ozs. (31.103 to 93.31 gms.)

---

**TRY MELLIN'S FOOD**  
**As a diet for Nursing Mothers.**



**Pepsinum—Pepsin.**

A proteolytic ferment obtained from the glandular layer of fresh stomachs from healthy pigs, and capable of digesting not less than 3,000 times its own weight of freshly coagulated and disintegrated egg albumen, when tested by the official process given.

DOSE: 1 to 3 grs. (0.0648 to 0.1944.)

**Pepsinum Saccharatum—Saccharated Pepsin.**

Should digest 300 times its weight of coagulated albumen.

DOSE: 5 to 30 grs. (0.324 to 1.944 gm.)

**Petrolatum Liquidum—Liquid Petrolatum.**

A mixture of hydrocarbons, chiefly of the marsh-gas series, obtained by distilling off the lighter and more volatile portions from petroleum, and purifying the residue when it has the desired consistence.

**Petrolatum Molle—Soft Petrolatum—Petrolatum, U. S. P., 1880. Soft Petroleum Ointment**

A mixture of hydrocarbons, chiefly of the marsh-gas series, obtained by distilling off the lighter and more volatile portions from petroleum, and purifying the residue when it has the desired melting point.

**Petrolatum Spissum—Hard Petrolatum—Petrolatum, U. S. P., 1880. Hard Petroleum Ointment.**

A mixture of hydrocarbons, chiefly of the marsh-gas series, obtained by distilling off the lighter and more volatile portions from petroleum, and purifying the residue when it has the desired melting point.

**Phosphorus—Phosphorus.**

DOSE: 1-100 to 1-20 gr. (0.00064 to 0.0032 gm.)

Preps: Oleum Phosphoratum. Pilulæ Phosphori. Spiritus Phosphori.

**Physostigma—Physostigma—Calabar Bean.**

The seed of *Physostigma venenosum*.

DOSE:  $\frac{1}{2}$  to 2 grs. (0.0324 to 0.1296 gm.)

Preps: Extractum Physostigmatis. Tinctura Physostigmatis.

**Physostigminæ Salicylas—Physostigmine Salicylate—Eserine Salicylate.**

The salicylate of an alkaloid obtained from *Physostigma*.

DOSE: 1-100 to 1-30 gr. (0.00064 to 0.00216 gm.)

**Physostigminæ Sulphas—Physostigmine Sulphate—Eserine Sulphate.**

The sulphate of an alkaloid obtained from *Physostigma*.

DOSE: 1-100 to 1-60 gr. (0.00064 to 0.0010 gm.)

**MELLIN'S FOOD**

Makes cow's milk easily digestible.

**Phytolaccae Fructus—*Phytolacca* Fruit—Phytolaccae Bacca, U. S. P., 1880. Poke Berry.**

The fruit of *Phytolacca decandra*.

DOSE: 5 to 30 grs. (0.324 to 1.944 gm.)

Prep: Extractum Phytolaccae Fluidum.

**Phytolaccae Radix—*Phytolacca* Root—Poke Root.**

The root of *Phytolacca decandra*.

DOSE: 5 to 30 grs. (0.324 to 1.944 gm.)

**Picrotoxinum—*Picrotoxin*.**

A neutral principle obtained from the seed of *Anamirta paniculata*.

DOSE: 1-120 to 1-60 gr. (0.00054 to 0.0010 gm.)

**Pilocarpinae Hydrochloras—*Pilocarpine* Hydrochlorate.**

The hydrochlorate of an alkaloid obtained from *Pilocarpus*.

DOSE:  $\frac{1}{4}$  to  $\frac{1}{2}$  gr. (0.0081 to 0.0324 gm.)

**Pilocarpus—*Pilocarpus*—Jaborandi.**

The leaflets of *Pilocarpus Selloanus* and of *Pilocarpus Jaborandi*.

DOSE: 5 to 60 grs. (0.324 to 3.888 gms.)

Prep: Extractum Pilocarpi Fluidum.

**Pilulae Aloes—*Pills of Aloes*.**

Purified Aloes, 0.13 gm.

DOSE: 2 to 5 pills.

**Pilulae Aloes et Asafoetidae—*Pill of Aloes and Asafoetida*.**

Purified Aloes, 0.09 gm. Asafoetida, 0.09 gm.

DOSE: 2 to 5 pills.

**Pilulae Aloes et Ferri—*Pills of Aloes and Iron*.**

Purified Aloes in fine powder, 0.07 gm. Dried Iron Sulphate, 0.07 gm.

DOSE: 1 to 3 pills.

**Pilulae Aloes et Mastiches—*Pills of Aloes and Mastic*.**

Purified Aloes, 0.13 gm. Mastic, 0.04 gm.

DOSE: 2 to 5 pills.

**Pilulae Aloes et Myrrhae—*Pills of Aloes and Myrrh*.**

Purified Aloes, 0.13 gm. Myrrh, 0.06 gm.

DOSE: 2 to 5 pills.

**Pilulae Antimonii Compositae—*Compound Pills of Antimony*—Plummer's Pills.**

DOSE: 1 to 3 pills.

**Pilulae Asafoetidae—*Pills of Asafoetida*.**

Asafoetida, 0.20 gm.

DOSE: 2 to 5 pills.

---

**MELLIN'S FOOD**  
Is readily assimilated.

**Pilulæ Catharticæ Compositæ—Compound Cathartic Pills.**

DOSE. 2 to 5 pills.

**Pilulæ Catharticæ Vegetabiles—Vegetable Cathartic Pills.**

DOSE 2 to 5 pills.

**Pilulæ Ferri Carbonatis—Pills of Carbonate of Iron—Ferruginous Pills. Chalybeate Pills. Bland's Pills.**

DOSE. 2 to 5 pills.

**Pilulæ Ferri Iodidi—Pills of Iodide of Iron.**

DOSE: 1 to 2 pills

**Pilulæ Opil—Pills of Opium.**

Powdered Opium, 0.065 gm.

DOSE: 1 to 2 pills.

**Pilulæ Phosphori—Pills of Phosphorus.**

Phosphorus, 0.0006 gm.

DOSE: 1 to 2 pills.

**Pilulæ Rhei—Pills of Rhubarb.**

Rhubarb, 0.20 gm.

DOSE: 1 to 5 pills.

**Pilulæ Rhei Compositæ—Compound Pills of Rhubarb.**

Rhubarb, 0.13 gm. Aloes, 0.10 gm. Myrrh, 0.06 gm.

DOSE: 1 to 3 pills.

**Pimenta—Pimenta—Allspice.**

The nearly ripe fruit of *Pimenta officinalis*.

DOSE: 5 to 30 grs. (0.324 to 1.944 gm.)

**Piper—Pepper—Black Pepper.**

Unripe fruit of *Piper nigrum*.

DOSE 5 to 20 grs (0.324 to 1.296 gm.)

Prep Oleoresina Piperis.

**Piperinum—Piperin.**

A neutral principle obtained from pepper and obtainable also from other plants of the natural order *Piperaceæ*

DOSE 1 to 10 grs (0.0648 to 0.648 gm.)

**Pix Burgundica—Burgundy Pitch.**

The prepared resinous exudation of *Abies excelsa*.

Preps Emplastrum Picis Burgundicæ. Emplastrum Picis Cantharidatum.

**Pix Liquida—Tar.**

An empyreumatic oleoresin obtained by the destructive distillation of the wood of *Pinus palustris* and of other species of *Pinus*.

DOSE:  $\frac{1}{4}$  to 1 dr (0.972 to 3.888 gm)

Preps Syrupus Picis Liquidæ Unguentum Picis Liquidæ.

**MELLIN'S FOOD**

Is a soluble dry extract from wheat and malt.

**Plumbi Acetas—Lead Acetate—Sugar of Lead.**

DOSE: 1 to 5 grs. (0.648 to 0.324 gm.)

Prep: Liquor Plumbi Subacetatis.

**Plumbi Carbonas Lead Carbonate—White Lead.**

Prep: Unguentum Plumbi Carbonatis.

**Plumbi Iodidum—Lead Iodide.**DOSE:  $\frac{1}{4}$  to 1 gr. (0.0162 to 0.0648 gm.)

Prep: Unguentum Plumbi Iodidi.

**Plumbi Nitras—Lead Nitrate.**DOSE:  $\frac{1}{4}$  to  $\frac{1}{2}$  gr. (0.0162 to 0.0324 gm.)**Plumbi Oxidum—Lead Oxide.**

Prep: Emplastrum Plumbi.

**Podophyllum—Podophyllum—May Apple.**The rhizome and roots of *Podophyllum peltatum*.

DOSE: 5 to 20 grs. (0.324 to 1.296 gm.)

Preps: Extractum Podophylli. Extractum Podophylli Fluidum. Resina Podophylli.

**Potassa—Potassa—Potassium Hydrate. Potassium Hydroxide. Caustic Potash.**

Preps: Liquor Potassie. Potassa cum Calce.

**Potassa cum Calce—Potassa with Lime.**

Potassa, 50%. Lime, 50%.

**Potassa Sulphurata—Sulphurated Potassa—Liver of Sulphur.**DOSE:  $\frac{1}{2}$  to 5 grs. (0.0324 to 0.324 gm.)**Potassii Acetas—Potassium Acetate.**

DOSE: 10 to 60 grs. (0.648 to 3.888 gms.)

**Potassii Bicarbonas—Potassium Bicarbonate.**

DOSE: 10 to 60 grs. (0.648 to 3.888 gm.)

**Potassii Bichromas—Potassium Bichromate.**

DOSE: 1-10 to 1 gr. (0.0064 to 0.0648 gm.)

**Potassii Bitartras—Potassium Bitartrate—Cream of Tartar.**

DOSE: 1 to 8 drs. (3.888 to 31.103 gm.)

Prep: Pulvis Jalapae Compositus.

**Potassii Bromidum—Potassium Bromide.**

DOSE: 10 to 60 grs. (0.648 to 3.888 gms.)

**Potassii Carbonas—Potassium Carbonate.**

DOSE: 10 to 30 grs. (0.648 to 1.944 gm.)

**Potassii Chloras—Potassium Chlorate.**

DOSE: 5 to 20 grs. (0.324 to 1.296 gm.)

Prep: Trochisci Potassi Chloratis.

**Potassii Citras—Potassium Citrate.**

DOSE: 15 to 60 grs. (0.972 to 3.888 gms.)

**Potassii Citras Effervescens—Effervescent Potassium Citrate.**

DOSE: 30 to 90 grs. (1.944 to 5.832 gms.)

---

**MELLIN'S FOOD****Is entirely free from starch and cane sugar.**

**Potassii Cyanidum—Potassium Cyanide.**DOSE:  $\frac{1}{8}$  to  $\frac{1}{4}$  gr. (0.0081 to 0.0162 gm.)**Potassii et Sodii Tartras—Potassium and Sodium Tartrate—Rochelle Salt.**DOSE: 1 to 2 grs.,  $\frac{1}{4}$  to 1 oz. (0.0648 to 0.1296, 7.776 to 31.103 gms.)**Potassii Ferrocyanidum—Potassium Ferrocyanide—Yellow Prussiate of Potash.**

DOSE: 10 to 30 grs. (0.648 to 1.944 gm.)

**Potassii Hypophosphis—Potassium Hypophosphite.**

DOSE: 5 to 30 grs. (0.324 to 1.944 gm.)

Prep: Syrupus Hypophosphitum.

**Potassii Iodidum—Potassium Iodide.**

DOSE: 5 to 60 grs. (0.324 to 3.888 gm.)

Prep: Unguentum Potassii Iodidi.

**Potassii Nitras—Potassium Nitrate—Saltpetre.**

DOSE: 10 to 30 grs. (0.648 to 1.944 gm.)

Prep: Argenti Nitras Dilutus. Charta Potassii Nitratis.

**Potassii Permanganas—Potassium Permanganate.**DOSE:  $\frac{1}{2}$  to 3 grs. (0.0324 to 0.1944 gm.)**Potassii Sulphas—Potassium Sulphate.**DOSE: 15 to 60 grs.,  $\frac{1}{2}$  to 4 drs. (0.972 to 3.888 gms, 1.944 to 15.552 gms.)**Prunum—Prune.**The fruit of *Prunus domestica*.

Prep: Confectio Sennæ.

**Prunus Virginiana—Wild Cherry.**The bark of *Prunus serotina*, collected in Autumn.DOSE:  $\frac{1}{2}$  to 1 dr. (1.944 to 3.888 gms.)

Preps: Extractum Pruni Virginianæ Fluidum. Infusum Pruni Virginianæ. Syrupus Pruni Virginianæ.

**Pulsatilla—Pulsatilla.**The herb of *Anemone Pulsatilla* and of *Anemone pratensis* collected soon after flowering.

DOSE: 1 to 5 grs. (0.0648 to 0.324 gm.)

**Pulvis Antimonialis—Antimonial Powder—James' Powder.**

Antimony Oxide, 33%. Precip. Calcium Phosphate, 67%.

DOSE: 3 to 10 grs. (0.1944 to 0.648 gm.)

**Pulvis Aromaticus—Aromatic Powder.**

DOSE: 10 to 30 grs. (0.648 to 1.944 gm.)

Prep: Extractum Aromaticum Fluidum.

**Pulvis Cretæ Compositus—Compound Chalk Powder.**

Prepared Chalk, 30%. Acacia, 20%. Sugar, 50%.

DOSE: 15 to 60 grs. (0.972 to 3.888 gm.)

Prep: Mistura Cretæ.

**MELLIN'S FOOD**

Stimulates the bowels to regular action.

**Pulvis Effervescens Compositus**—*Compound Effervescing Powder*—Seidlitz Powder.

DOSE: 1 powder.

**Pulvis Glycyrrhizæ Compositum**—*Compound Powder of Glycyrrhiza*.

DOSE:  $\frac{1}{2}$  to 2 drs. (1.944 to 7.776 gms.)

**Pulvis Ipecacuanhæ et Opil**—*Powder of Ipecac and Opium*—Dover's Powder.

Ipecac, 10%. Opium, 10%.

DOSE: 5 to 15 grs. (0.324 to 0.972 gm.)

**Pulvis Jalapæ Compositus**—*Compound Powder of Jalap*—Pulvis Purgans.

Jalap, 35%.

DOSE.  $\frac{1}{4}$  to 1 dr. (0.972 to 3.888 gms.)

**Pulvis Morphinæ Compositus**—*Compound Powder of Morphine*—Tully's Powder.

DOSE: 5 to 15 grs. (0.324 to 0.972 gm.)

**Pulvis Rhei Compositus**—*Compound Powder of Rhubarb*.

Rhubarb, 25%.

DOSE: 1 to 3 drs. (3.888 to 11.664 gms.)

**Pyrethrum**—*Pyrethrum*—Pellitory.

The root of *Anacyclus Pyrethrum*.

DOSE:  $\frac{1}{2}$  to 1 dr. (1.944 to 3.888 gms.)

Prep: *Tinctura Pyrethri*.

**Pyrogallol**—*Pyrogallol*—Pyrogallic Acid.

A triatomic phenol, obtained chiefly by the dry distillation of gallic acid.

**Pyroxylinum**—*Pyroxylin*—Soluble Gun Cotton. Colloxylin.

Prep: *Collodium*.

**Quassia**—*Quassia*.

The wood of *Pieræna excelsa*.

Preps. *Extractum Quassiae*. *Extractum Quassiae Fluidum*.

*Tinctura Quassiae*.

**Quercus Alba**—*White Oak*.

The bark of *Quercus alba*.

DOSE:  $\frac{1}{4}$  to 1 dr. (0.972 to 3.888 gms.)

**Quillaja**—*Quillaja*—*Quillaia*, U. S. P. 1880. Soap Bark.

The inner bark of *Quillaja Saponaria*.

DOSE:  $\frac{1}{4}$  to  $\frac{1}{2}$  dr. (0.972 to 1.944 gm.)

Prep: *Tinctura Quillajæ*.

---

"With children sick with Typhoid Fever, I have found MELLIN'S an excellent food."

**Quinidinæ Sulphas—Quinidine Sulphate.**

The neutral sulphate of an alkaloid obtained from the bark of several species of *Cinchona*.

DOSE 1 to 30 grs. (0.0648 to 1.944 gm.)

**Quinina—Quinine.**

An alkaloid obtained from the bark of various species of *Cinchona*.

DOSE: 1 to 30 grs. (0.0648 to 1.944 gm.)

Preps: Ferri et Quininæ Citras. Ferri et Quininæ Citras Solubilis.

**Quininæ Bisulphas—Quinine Bisulphate.**

DOSE. 1 to 30 grs. 0.0648 to 1.944 gm.)

**Quininæ Hydrobromas—Quinine Hydrobromate.**

DOSE: 1 to 30 grs. (0.0648 to 1.944 gm.)

**Quininæ Hydrochloras—Quinine Hydrochlorate.**

DOSE 1 to 30 grs. (0.0648 to 1.944 gm.)

**Quininæ Sulphas—Quinine Sulphate.**

DOSE: 1 to 30 grs. (0.0648 to 1.944 gm.)

Prep. Syrupus Ferri, Quininæ et Strychninæ Phosphatum.

**Quininæ Valerianas—Quinine Valerianate.**

DOSE 1 to 3 grs. (0.0648 to 0.1944 gm.)

**Resina—Resin—Colophony.**

The residue left after distilling off the volatile oil from Turpentine.

Preps: Ceratum Resinæ. Emplastrum Resinæ.

**Resina Copaibæ—Resin of Copaiba.**

The residue left after distilling off the volatile oil from Copaiba.

DOSE: 5 to 20 grs. (0.324 to 1.296 gm.)

**Resina Jalapæ—Resin of Jalap.**

DOSE: 1 to 5 grs. (0.0648 to 0.324 gm.)

**Resina Podophylli—Resin of Podophyllum.**

DOSE:  $\frac{1}{4}$  to 2 grs. (0.0081 to 0.1296 gm.)

Prep: Pilulæ Catharticæ Vegetabiles.

**Resina Scammonii—Resin of Scammony.**

DOSE: 2 to 8 grs. (0.1296 to 0.5184 gm.)

**Resorcinum—Resorcin—Resorcinol. Metadi-oxybenzol.**

A diatomic phenol.

DOSE: 3 to 10 grs. (0.1944 to 0.648 gm.)

**Rhamnus Purshiana—Cascara Sagrada.**

The bark of *Rhamnus purshiana*.

DOSE.  $\frac{1}{4}$  to 1 dr. (0.972 to 3.888 gms.)

Prep: Extractum Rhamni Purshianæ Fluidum.

**MELLIN'S FOOD**

Contains the nitrogenous and phosphatic matters essential for the growing child.

**Rheum—Rhubarb.**

The root of *Rheum officinale*.

DOSE: 5 to 30 grs. (0.324 to 1.944 gm.)

Preps: Extractum Rhei. Extractum Rhei Fluidum. Pilulæ Rhei. Pilulæ Rhei Compositæ. Pulvis Rhei Compositus. Tinctura Rhei. Tinctura Rhei Aromatica. Tinctura Rhei Dulcis.

**Rhus Glabra—Rhus Glabra.**

The fruit of *Rhus glabra*.

DOSE:  $\frac{1}{4}$  to 1 dr. (0.972 to 3.888 gms.)

Prep: Extractum Rhois Glabræ Fluidum.

**Rhus Toxicodendron—Rhus Toxicodendron—Poison Ivy.**

The fresh leaves of *Rhus radicans*.

DOSE: 1 to 5 grs. (0.0648 to 0.324 gm.)

**Rosa Centifolia—Pale Rose.**

The petals of *Rosa centifolia*.

Preps: Syrupus Sarsaparillæ Compositus.

**Rosa Gallica—Red Rose.**

The petals of *Rosa gallica*, collected before expanding.

Preps: Confectio Rosæ. Extractum Rosæ Fluidum. Pilulæ Aloes et Mastiches.

**Rubus—Rubus—Blackberry.**

The bark of the root of *Rubus villosus*, *Rubus canadensis* and *Rubus trivialis*.

Prep: Extractum Rubi Fluidum.

**Rubus Idæus—Raspberry.**

The fruit of *Rubus idæus*.

Prep: Syrupus Rubi Idæi.

**Rumex—Rumex—Yellow Dock.**

The root of *Rumex crispus* Linne, and of some other species of *Rumex*.

DOSE:  $\frac{1}{4}$  to 1 dr. (0.972 to 3.888 gms.)

Prep: Extractum Rumicis Fluidum.

**Sabina—Savine.**

The tops of *Juniperus Sabina*.

DOSE: 5 to 15 grs. (0.324 to 0.972 gm.)

Prep: Extractum Sabinæ Fluidum.

**Saccharum—Sugar.**

The refined sugar obtained from *Saccharum officinarum* and from various species and varieties of *Sorghum* also from one or more varieties of *Beta vulgaris*.

Preps: Syrupus. (Compound Syrups, etc.)

**Saccharum Lactis—Sugar of Milk.**

A peculiar crystalline sugar, obtained from the whey of cow's milk by evaporation, and purified by recrystallization.

**MELLIN'S FOOD**

Prepared with milk fulfils every requirement of a food for infants.



**Salicinum—Salicin.**

A neutral principle obtained from several species of *Salix* and *Populus*.

DOSE. 10 to 30 grs. (0.648 to 1.944 gm.)

**Salol—Salol—Phenyl Salicylate.**

The salicylic ether of phenol.

DOSE: 3 to 15 grs. (0.1944 to 0.972 gm.)

**Salvia—Salvia—Sage.**

The leaves of *Salvia officinalis*.

DOSE:  $\frac{1}{2}$  to 1 dr. (0.972 to 3.888 gms.)

**Sambucus—Sambucus—Elder.**

The flowers of *Sambucus canadensis*.

DOSE:  $\frac{1}{2}$  to 1 dr. (1.944 to 3.888 gms.)

**Sanguinaria—Sanguinaria—Bloodroot.**

The rhizome of *Sanguinaria canadensis* collected in Autumn.

DOSE. 2 to 30 grs. (0.1296 to 1.944 gm.)

Preps. Extractum Sanguinarum Fluidum. Tinctura Sanguinarum.

**Santalum Rubrum—Red Saunders.**

The wood of *Pterocarpus santalinus*.

**Santonica—Santonica—Levant Wormseed.**

The unexpanded flower heads of *Artemisia pauciflora*.

DOSE: 3 to 15 grs. (0.1944 to 0.972 gm.)

**Santoninum—Santonin.**

A neutral principle obtained from Santonica.

DOSE. 1 to 3 grs. (0.0648 to 0.1944 gm.)

Prep: Trochisci Santonini.

**Sapo—Soap—White Castile Soap.**

Soap prepared from soda and olive oil.

DOSE. 5 to 30 grs. (0.324 to 1.944 gm.)

Preps: Emplastrum Saponis. Linimentum Saponis.

**Sapo Mollis—Soft Soap—Sapo Viridis, U. S. P., 1880. Green Soap.**

Prep: Linimentum Saponis Mollis.

**Sarsaparilla—Sarsaparilla.**

The root of *Smilax officinalis*, *Smilax medica*, *Smilax papyracea* and of other undetermined species of *Smilax*.

DOSE:  $\frac{1}{2}$  to 1 dr. (1.944 to 3.888 gms.)

Preps: Decoctum Sarsaparillae Compositum. Extractum Sarsaparillae Fluidum. Extractum Sarsaparillae Fluidum. Compositum.

**Sassafras—Sassafras.**

The bark of the root of *Sassafras variifolium*.

DOSE.  $\frac{1}{2}$  to 1 dr. (0.972 to 3.888 gms.)

**Sassafras Medulla—Sassafras Pith.**

The pith of *Sassafras variifolium*.

Prep Mucilago Sassafras Medullae.

**MELLIN'S FOOD**

Contains a large amount of proteids and soluble phosphates.

**Scammonium—Scammony.**

Resinous exudate from living root of *Convolvulus Scammonia*.

DOSE: 5 to 15 grs. (0.324 to 0.972 gm.)

Prep: Resina Scammonii.

**Scilla—Squill.**

The bulb of *Urginea maritima*, deprived of its dry membranaceous outer scales and cut into thin slices, the central portions being rejected.

DOSE: 1 to 3 grs. (0.0648 to 0.1944 gm.)

Preps: Acetum Scillæ. Extractum Scillæ Fluidum. Tinctura Scillæ.

**Scoparius—Scoparius—Broom.**

The tops of *Cytisus Scoparius*.

DOSE:  $\frac{1}{2}$  to 1 dr. (0.972 to 3.888 gms.)

Prep: Extractum Scoparii Fluidum.

**Scutellaria—Scutellaria—Scullcap.**

The herb of *Scutellaria lateriflora*.

DOSE: 1 to 2 drs. (3.888 to 7.776 gms.)

Prep: Extractum Scutellariæ Fluidum.

**Senega—Senega.**

The root of *Polygala Senega*.

DOSE: 10 to 20 grs. (0.648 to 1.296 gm.)

Prep: Extractum Senegæ Fluidum.

**Senna—Senna.**

The leaflets of *Cassia angustifolia*.

DOSE: 15 grs. to 2 drs. (0.972 to 7.776 gms.)

Preps: Confectio Sennæ. Extractum Sennæ Fluidum. Infusum Sennæ Compositum. Pulvis Glycyrrhizæ Compositus. Syrupus Sennæ.

**Serpentaria—Serpentaria—Virginia Snakeroot.**

The rhizome and roots of *Aristolochia Serpentaria* and of *Aristolochia reticulata*.

DOSE: 10 to 30 grs. (0.648 to 1.944 gm.)

Preps: Extractum Serpentariæ Fluidum. Tinctura Cinchonæ Composita. Tinctura Serpentariæ.

**Sevum—Suet—Mutton Suet.**

The internal fat of the abdomen of *Ovis Arles* purified by melting and straining.

**Sinapis Alba—White Mustard.**

The seed of *Brassica alba*.

DOSE: 1 to 4 drs. (3.888 to 15.552 gms.)

**Sinapis Nigra—Black Mustard.**

The seed of *Brassica nigra*.

DOSE: 1 to 4 drs. (3.888 to 15.552 gms.)

Prep: Charta Sinapis.

**Soda—Soda—Sodium Hydrate. Sodium Hydroxide. Caustic Soda.**

Prep: Liquor Sodæ.

**MELLIN'S FOOD**

Actually assists to digest milk or other foods with which it is mixed.

**Sodii Acetas—Sodium Acetate.**

DOSE 10 to 40 grs. (0.648 to 2.592 gms.)

**Sodii Arsenas—Sodium Arsenate—Sodii Arsenias, U. S. P., 1880.**

DOSE: 1/24 to 1/4 gr (0.0027 to 0.0081 gm.)

Prep Liquor Sodii Arsenatis.

**Sodii Benzoas—Sodium Benzoate.**

DOSE. 10 to 60 grs (0.648 to 3.888 gms.)

**Sodii Bicarbonas—Sodium Bicarbonate**

DOSE: 10 to 60 grs (0.648 to 3.888 gms.)

Preps Mistura Rhei et Sodæ. Trochisci Sodii Bicarbonatis.

**Sodii Bisulphis—Sodium Bisulphite.**

DOSE: 10 to 30 grs. (0.648 to 1.944 gm.)

**Sodii Boras—Sodium Borate—Borax.**

DOSE: 5 to 30 grs. (0.324 to 1.944 gm.)

**Sodii Bromidum—Sodium Bromide.**

DOSE 10 to 60 grs. (0.648 to 3.888 gms.)

**Sodii Carbonas—Sodium Carbonate.**

DOSE. 5 to 20 grs (0.324 to 1.296 gm.)

Prep: Sodii Carbonas Exsiccatus.

**Sodii Carbonas Exsiccatus—Dried Sodium Carbonate**

DOSE: 5 to 15 grs. (0.324 to 0.972 gm.)

**Sodii Chloras—Sodium Chlorate.**

DOSE 3 to 15 grs. (0.1944 to 0.972 gm.)

**Sodii Chloridum—Sodium Chloride.**

DOSE 10 to 60 grs. (0.648 to 3.888 gms.)

**Sodii Hypophosphis—Sodium Hypophosphite.**

DOSE. 5 to 20 grs. (0.324 to 1.296 gm.)

Prep: Syrupus Hypophosphitum.

**Sodii Hyposulphis—Sodium Hyposulphite—Sodium Thiosulphate.**

DOSE 5 to 24 grs (0.324 to 1.555 gm.)

**Sodii Iodidum—Sodium Iodide.**

DOSE: 5 to 60 grs. (0.324 to 3.888 gms.)

**Sodii Nitras—Sodium Nitrate.**

DOSE. 8 to 40 grs. (0.5184 to 2.592 gm.)

**Sodii Nitris—Sodium Nitrite.**

DOSE 1 to 3 grs (0.0648 to 0.1944 gm.)

Prep Spiritus Ætheris Nitrosi.

**Sodii Phosphas—Sodium Phosphate—Sodium Orthophosphate.**

DOSE: 1 to 8 drs (3.888 to 31.103 gms.)

**Sodii Pyrophosphas—Sodium Pyrophosphate.****Sodii Salicylas—Sodium Salicylate.**

DOSE 8 to 60 grs (0.5184 to 3.888 gms.)

---

**MELLIN'S FOOD**

proved its greater nutritive value over all other artificial foods.

**Sodii Sulphas**—*Sodium Sulphate*—Glauber's Salt.

DOSE: 1 to 8 drs. (3.888 to 31.103 gms.)

**Sodii Sulphis**—*Sodium Sulphite*.

DOSE: 10 to 60 grs. (0.648 to 3.888 gms.)

**Sodii Sulphocarbolas**—*Sodium Sulphocarbolate*—Sodium Paraphenolsulphonate.

DOSE: 8 to 30 grs. (0.5184 to 1.944 gm.)

**Sparteinae Sulphas**—*Sparteine Sulphate*.

The neutral sulphate of an alkaloid obtained from *Scoparius*.

DOSE:  $\frac{3}{4}$  to 1 gr. (0.0081 to 0.0648 gm.)

**Spigelia**—*Spigelia*—Pinkroot.

The rhizome and roots of *Spigelia marilandica*.

DOSE:  $\frac{1}{4}$  to 2 drs. (0.972 to 7.776 gms.)

Prep: *Extractum Spigeliae Fluidum*.

**Spiritus Aetheris**—*Spirit of Ether*.

Ether, 32.5%. Alcohol, 67.5%.

DOSE: 15 to 20 min. (0.92 to 1.23 c. c.)

**Spiritus Aetheris Compositus**—*Compound Spirit of Ether*—Hoffman's Anodyne.

Ether, 32.5%. Alcohol, 68%. Ethereal Oil, 2.5%.

DOSE: 15 to 60 min. (0.92 to 3.70 c. c.)

**Spiritus Aetheris Nitrosi**—*Spirit of Nitrous Ether*.

An alcoholic solution of Ethyl Nitrite, yielding when freshly prepared and tested in a nitrometer, not less than 11 times its own volume of Nitrogen Dioxide.

DOSE:  $\frac{1}{4}$ , 1 to 2 fl. drs. (0.92, 3.70 to 7.39 c. c.)

**Spiritus Ammoniae**—*Spirit of Ammonia*.

An alcoholic solution of Ammonia, containing 10% by weight of the gas.

DOSE: 10 to 30 min. (0.62 to 1.85 c. c.)

**Spiritus Ammoniae Aromaticus**—*Aromatic Spirit of Ammonia*.

DOSE: 30 to 60 min. (1.85 to 3.70 c. c.)

Preps: *Tinctura Guaiaci Ammoniata*. *Tinctura Valerianae Ammoniata*.

**Spiritus Amygdalae Amaræ**—*Spirit of Bitter Almond*—Essence of Bitter Almond.

Oil of Bitter Almond, 1%.

**Spiritus Anisi**—*Spirit of Anise*.

Oil of Anise, 10%.

DOSE: 1 to 2 fl. drs. (3.70 to 7.39 c. c.)

**Spiritus Aurantii**—*Spirit of Orange*.

Oil of Orange Peel, 5%.

## MELLIN'S FOOD

Is a most useful addition to the diet of healthy children.

- Spiritus Aurantii Compositus**—*Compound Spirit of Orange.*  
 Prep Elixir Aromaticum.
- Spiritus Camphoræ**—*Spirit of Camphor.*  
 Camphor, 10%.  
 DOSE: 5 to 40 min. (0.31 to 2.46 c. c.)
- Spiritus Chloroformi**—*Spirit of Chloroform.*  
 Chloroform, 6%.  
 DOSE. 10 to 60 min. (0.62 to 3.70 c. c.)
- Spiritus Cinnamomi**—*Spirit of Cinnamon.*  
 Oil of Cinnamon, 10%.  
 DOSE. 10 to 20 min (0.62 to 1.23 c. c.)
- Spiritus Frumenti**—*Whiskey.*  
 An alcoholic liquid obtained by the distillation of the mash of fermented grain (usually of mixtures of corn, wheat and rye), and at least two years old.
- Spiritus Gaultheriæ**—*Spirit of Gaultheria.*  
 Oil of Gaultheria, 5%.  
 DOSE: 10 to 30 min. (0.62 to 1.85 c. c.)
- Spiritus Glonoini**—*Spirit of Glonoin*—*Spirit of Nitroglycerin.*  
 An alcoholic solution of Glonoin, containing 1% by weight of the substance.  
 DOSE. 1 min. (0.06 c. c.)
- Spiritus Juniperi**—*Spirit of Juniper.*  
 Oil of Juniper, 5%.  
 DOSE: 30 to 60 min. (1.85 to 3.70 c. c.)
- Spiritus Juniperi Compositus**—*Compound Spirit of Juniper.*  
 DOSE 1 to 4 fl. drs. (3.70 to 14.79 c. c.)
- Spiritus Lavandulæ**—*Spirit of Lavender.*  
 Oil of Lavender Flowers, 5%.  
 DOSE. 15 to 60 min. (0.92 to 3.70 c. c.)
- Spiritus Limonis**—*Spirit of Lemon*—*Essence of Lemon.*  
 Oil of Lemon, 5%.
- Spiritus Menthæ Piperitæ**—*Spirit of Peppermint*—*Essence of Peppermint.*  
 Oil of Peppermint, 10%.  
 DOSE: 10 to 30 min. (0.62 to 1.85 c. c.)  
 Prep: Mistura Rhei et Sodæ.
- Spiritus Menthæ Viridis**—*Spirit of Spearmint*—*Essence of Spearmint.*  
 Oil of Spearmint, 10%.  
 DOSE. 15 to 40 min. (0.92 to 2.46 c. c.)
- Spiritus Myrciæ**—*Spirit of Myrcia*—*Bay Rum.*  
 Oil of Myrcia, 0.8%.

---

## MELLIN'S FOOD

Is unrivalled as an artificial food for children.

**Spiritus Myristicæ—Spirit of Nutmeg—Es-  
sence of Nutmeg.**

Oil of Nutmeg, 5%.

DOSE 15 to 60 min. (0.92 to 3.70 c. c.)

**Spiritus Phosphori—Spirit of Phosphorus—  
Tincture of Phosphorus.**

Phosphorus, 0.12%.

Prep: Elixir Phosphori.

**Spiritus Vini Gallici—Brandy.**

An alcoholic liquid, obtained by the distillation of the fermented, unmodified juice of fresh grapes, and at least four years old.

**Staphisagria—Staphisagria—Stavesacre.**

The seed of *Delphinium Staphisagria*.

**Stillingia—Stillingia—Queen's Root.**

The root of *Stillingia sylvatica*.

DOSE:  $\frac{1}{4}$  to 1 dr. (0.972 to 3.888 gms.)

Prep Extractum Stillingiæ Fluidum.

**Stramonii Folia—Stramonium Leaves.**

The leaves of *Datura Stramonium*.

DOSE 1 to 5 grs. (0.0648 to 0.324 gm.)

**Stramonii Semen—Stramonium Seed.**

The seed of *Datura Stramonium*.

DOSE 1 to 3 grs. (0.0648 to 0.1944 gm.)

Preps Extractum Stramonii Seminis. Extractum Stramonii Seminis Fluidum. Tinctura Stramonii Seminis.

**Strontii Bromidum—Strontium Bromide.**

DOSE: 5 to 10 grs. (0.324 to 0.648 gm.)

**Strontii Iodidum—Strontium Iodide.**

DOSE 5 to 10 grs. (0.324 to 0.648 gm.)

**Strontii Lactas—Strontium Lactate.**

DOSE. 5 to 10 grs. (0.324 to 0.648 gm.)

**Strophanthus—Strophanthus.**

The seed of *Strophanthus hispidus* deprived of its long awn.

Prep. Tinctura Strophanthi.

**Strychnina—Strychnine.**

An alkaloid obtained from *Nux Vomica*, and also obtainable from other plants of the natural order *Loganiaceæ*.

DOSE 1-60 to 1-20 gr. (0.001 to 0.0032 gm.)

Preps Ferri et Strychninæ Citras. Syrupus Ferri, Quininae et Strychninæ Phosphatum.

**Strychninæ Sulphas—Strychnine Sulphate.**

DOSE: 1-32 to 1-12 gr. (0.0020 to 0.0054 gm.)

**Styrax—Storax.**

Prepared from the inner bark of *Liquidambar orientalis*.

DOSE: 5 to 20 grs. (0.324 to 1.296 gm.)

Prep: Tinctura Benzoini Composita.

---

**MELLIN'S FOOD**  
Is a true Liebig's Food.

**Sulphuris Iodidum—Sulphur Iodide**

Washed Sulphur, 20%. Iodine, 80%.

DOSE: 1 to 4 grs. (0.0648 to 0.2592 gm.)

**Sulphur Lotum—Washed Sulphur.**

DOSE: 15 to 90 grs. (0.972 to 5.832 gms.)

Preps: Pulvis Glycyrrhizæ Compositus. Unguentum Sulphuris.

**Sulphur Præcipitatum—Precipitated Sulphur.**

DOSE: 15 to 90 grs. (0.972 to 5.832 gms.)

**Sulphur Sublimatum—Sublimed Sulphur.**

DOSE: 1 to 3 drs. (3.888 to 11.664 gms.)

Preps. Sulphur Lotum. Sulphur præcipitatum.

**Sumbul—Sumbul.**The root of *Ferula Sumbul*.

DOSE: 30 to 60 grs. (1.944 to 3.888 gms.)

Prep: Tinctura Sumbul.

**Suppositoria—Suppositories.**

Medicinal Ingredient, the prescribed quantity, Oil of Theobroma, a sufficient quantity.

**Suppositoria Glycerini—Suppositories of Glycerin.****Syrupus—Syrup.**

Saturated Solution of Sugar.

**Syrupus Acaciæ—Syrup of Acacia.**

Mucilage of Acacia, 25%.

DOSE: 1 to 8 fl. drs. (3.70 to 29.57 c. c.)

**Syrupus Acidi Citrici—Syrup of Citric Acid.**

Citric Acid, 1%.

DOSE:  $\frac{3}{4}$  to 1 fl. oz. (7.39 to 29.57 c. c.)**Syrupus Acidi Hydriodici—Syrup of Hydriodic Acid.**

A syrupy liquid containing about 1% by weight of absolute Hydriodic Acid, or about 1.3 gm. in 100 c. c.

DOSE: 30 to 60 min. (1.85 to 3.70 c. c.)

**Syrupus Allii—Syrup of Garlic.**

DOSE: 1 to 2 fl. drs. (3.70 to 7.39 c. c.)

**Syrupus Althææ—Syrup of Althæa.**

DOSE: Freely.

**Syrupus Amygdalæ—Syrup of Almond.**DOSE:  $\frac{3}{4}$  to 1 fl. oz. (7.39 to 29.57 c. c.)**Syrupus Aurantii—Syrup of Orange.**DOSE:  $\frac{3}{4}$  to 1 fl. oz. (7.39 to 29.57 c. c.)**Syrupus Aurantii Florum—Syrup of Orange Flowers.**

DOSE: 1 to 2 fl. drs. (3.70 to 7.39 c. c.)

**MELLIN'S FOOD**

Is most valuable as a nutritious food.

**Syrupus Calcii Lactophosphatis—Syrup of Calcium Lactophosphate.**

DOSE. 1 to 2 fl. drs. (3.70 to 7.39 c. c.)

**Syrupus Calcis—Syrup of Lime.**DOSE:  $\frac{1}{2}$  to 1 fl. dr. (1.85 to 3.70 c. c.)**Syrupus Ferri Iodidi—Syrup of Ferrous Iodide.**

A syrupy liquid containing about 10% by weight of Ferrous Iodide, or about 13.4 gm. in 100 c. c.

DOSE. 15 to 30 min. (0.92 to 1.85 c. c.)

**Syrupus Ferri, Quininæ et Strychninæ Phosphatum—Syrup of the Phosphates of Iron, Quinine and Strychnine.**DOSE:  $\frac{1}{2}$  to 1 fl. dr. (1.85 to 3.70 c. c.)**Syrupus Hypophosphitum—Syrup of Hypophosphites.**

DOSE: 1 to 2 fl. drs. (3.70 to 7.39 c. c.)

Prep: Syrupus Hypophosphitum cum Ferro.

**Syrupus Hypophosphitum cum Ferro—Syrup of Hypophosphites with Iron.**DOSE:  $\frac{1}{2}$  to 1  $\frac{1}{2}$  fl. drs. (1.85 to 5.54 c. c.)**Syrupus Ipecacuanhæ—Syrup of Ipecac.**

Fluid Extract of Ipecac, 7%.

DOSE: 5 to 30 min.; 2 to 6 fl. drs. (0.31 to 1.85 c. c.; 7.39 to 22.18 c. c.)

**Syrupus Kramerizæ—Syrup of Krameria**

Fluid Extract of Krameria, 45%.

DOSE: 2 to 6 fl. drs. (7.39 to 22.18 c. c.)

**Syrupus Lactucarii—Syrup of Lactucarium.**

Tincture of Lactucarium, 10%.

DOSE:  $\frac{1}{2}$  to 2 fl. drs. (1.85 to 7.39 c. c.)**Syrupus Picis Liquidæ—Syrup of Tar.**

DOSE: 1 to 4 fl. drs. (3.70 to 14.79 c. c.)

**Syrupus Pruni Virginianæ—Syrup of Wild Cherry.**

DOSE: 1 to 4 fl. drs. (3.70 to 14.79 c. c.)

**Syrupus Rhei—Syrup of Rhubarb.**

Fluid Extract of Rhubarb, 10%.

DOSE: 2 to 6 fl. drs. (7.39 to 22.18 c. c.)

**Syrupus Rhei Aromaticus—Aromatic Syrup of Rhubarb.**

Aromatic Tincture of Rhubarb, 15%.

DOSE: 2 to 6 fl. drs. (7.39 to 22.18 c. c.)

**Syrupus Rosæ—Syrup of Rose.**

Fluid Extract of Rose, 12.5%.

DOSE: 1 fl. dr. (3.70 c. c.)

---

**TRY MELLIN'S FOOD**  
**For Hand-fed Infants.**



**Syrupus Rubi—Syrup of Rubus.**

Fluid Extract of Rubus, 25%.

DOSE: 1 to 2 fl. drs. (3.70 to 7.39 c. c.)

**Syrupus Rubi Idæi—Syrup of Raspberry.**

DOSE: 1 to 3 fl. drs. (3.70 to 11.09 c. c.)

**Syrupus Sarsaparillæ Compositus—Compound Syrup of Sarsaparilla.**

Fluid Extract of Sarsaparilla, 20%. Fluid Extracts of Glycyrrhiza and Senna, each 1.5%.

DOSE: 2 to 4 fl. drs. (7.39 to 14.79 c. c.)

**Syrupus Scillæ—Syrup of Squill.**

Vinegar of Squill, 45%.

DOSE:  $\frac{1}{2}$  to 1 fl. dr. (1.85 to 3.70 c. c.)**Syrupus Scillæ Compositus—Compound Syrup of Squill.**

Fluid Extract of Squill, 8%. Fluid Extract Senega 8%. Antimony and Potassium Tartrate, 0.2 gm. in 100 c. c.

DOSE: 15 to 60 min.; 1 to 2 fl. drs. (0.92 to 3.70 c. c.; 3.70 to 7.39 c. c.)

**Syrupus Senegæ—Syrup of Senega.**

Fluid Extract of Senega, 20%.

DOSE:  $\frac{1}{2}$  to 2 fl. drs. (1.85 to 7.39 c. c.)**Syrupus Sennæ—Syrup of Senna.**

DOSE: 2 to 4 fl. drs. (7.39 to 14.79 c. c.)

**Syrupus Tolutanus—Syrup of Tolu.**

DOSE: 2 to 6 fl. drs. (7.39 to 22.18 c. c.)

**Syrupus Zingiberis—Syrup of Ginger.**

Fluid Extract of Ginger, 3%.

DOSE: 2 to 6 fl. drs. (7.39 to 22.18 c. c.)

**Tabacum—Tobacco.**The commercial dried leaves of *Nicotiana Tabacum*.

DOSE: 2 to 5 grs. (0.129 to 0.324 gm.)

**Tamarindus—Tamarind.**The preserved pulp of the fruit of *Tamarindus indica*.

DOSE: 1 dr. to 1 oz. (3.888 to 31.163 gm.)

Prep: Confectio Sennæ.

**Tanacetum—Tansy.**The leaves and flowering tops of *Tanacetum vulgare*.

DOSE: 30 to 60 grs. (1.944 to 3.888 gms.)

**Taraxacum—Taraxacum—Dandelion.**The root of *Taraxacum officinale* gathered in Autumn.

DOSE: 1 to 4 drs. (3.888 to 15.552 gms.)

Preps: Extractum Taraxaci. Extractum Taraxaci Fluidum.

**Terebenum—Terebene.**

A liquid consisting chiefly of Pinene, and containing not more than very small proportions of Terpinene and Dipentene.

DOSE: 3 to 10 min. (0.18 to 0.62 c. c.)

**TRY MELLIN'S FOOD**  
For Invalids.

**Terebinthina—Turpentine.**

A concrete oleoresin obtained from *Pinus palustris* and from other species of *Pinus*.

DOSE: 5 to 30 min. (0.31 to 1.85 c. c.)

**Terebinthina Canadensis—Canada Turpentine—Canada Balsam. Balsam of Fir.**

A liquid oleoresin obtained from *Abies balsamea*.

DOSE 5 to 30 min. (0.31 to 1.85 c. c.)

**Terpini Hydras—Terpin Hydrate.**

The hydrate of the diatomic alcohol Terpin.

DOSE 2 to 10 grs. (0.1296 to 0.648 gm.)

**Thymol—Thymol.**

A phenol occurring in the volatile oils of *Thymus vulgaris*, *Monarda punctata* and *Carum Ajowan*.

DOSE:  $\frac{1}{2}$  to 2 grs. (0.0324 to 0.1296 gm.)

**Tinctura Aconiti—Tincture of Aconite.**

DOSE: 1 to 3 min. (0.06 to 0.18 c. c.)

**Tinctura Aloes—Tincture of Aloes.**

DOSE: 5 to 10 min.;  $\frac{1}{2}$  to 4 fl. drs. (0.31 to 0.62 c. c., 1.85 to 14.79 c. c.)

**Tinctura Aloes et Myrrhæ—Tincture of Aloes and Myrrh.**

DOSE:  $\frac{1}{2}$  to 2 fl. drs. (1.85 to 7.39 c. c.)

**Tinctura Arnicæ Florum—Tincture of Arnica Flowers.**

DOSE: 10 to 30 min. (0.62 to 1.85 c. c.)

**Tinctura Arnicæ Radicis—Tincture of Arnica Root.**

DOSE. 20 to 30 min. (1.23 to 1.85 c. c.)

**Tinctura Asafœtidæ—Tincture of Asafetida.**

DOSE: 10 to 40 min. (0.62 to 2.46 c. c.)

**Tinctura Aurantii Amari—Tincture of Bitter Orange Peel.**

DOSE. 1 to 2 fl. drs. (3.70 to 7.39 c. c.)

**Tinctura Aurantii Dulcis—Tincture of Sweet Orange Peel.**

Flavoring.

**Tinctura Belladonnæ Foliorum—Tincture of Belladonna Leaves—Tinctura Belladonnæ, Pharm. 1880.**

DOSE: 5 to 15 min. (0.31 to 0.92 c. c.)

**Tinctura Benzoini—Tincture of Benzoin.**

DOSE: 10 to 40 min. (0.62 to 2.46 c. c.)

**Tinctura Benzoini Composita—Compound Tincture of Benzoin.**

DOSE: 15 to 60 min. (0.92 to 3.70 c. c.)

---

**TRY MELLIN'S FOOD**  
**In Typhoid Fever.**

**Tinctura Bryoniæ—Tincture of Bryonia.**

DOSE: 1 to 2 fl. drs. (3.70 to 7.39 c. c.)

**Tinctura Calendulæ—Tincture of Calendula.**

DOSE: 1 to 4 fl. drs. (3.70 to 14.79 c. c.)

**Tinctura Calumbæ—Tincture of Calumba.**

DOSE: 1 to 4 fl. drs. (3.70 to 14.79 c. c.)

**Tinctura Cannabis Indicæ—Tincture of Indian Cannabis.**

DOSE: 5 to 20 min. (0.31 to 1.23 c. c.)

**Tinctura Cantharidis—Tincture of Cantharides.**

DOSE: 3 to 10 min. (0.18 to 0.62 c. c.)

**Tinctura Capsici—Tincture of Capsicum.**

DOSE: 15 to 60 min. (0.92 to 3.70 c. c.)

**Tinctura Cardamomi—Tincture of Cardamom.**

DOSE: 1 to 2 fl. drs. (3.70 to 7.39 c. c.)

**Tinctura Cardamomi Composita—Compound Tincture of Cardamom.**

DOSE: 1 to 3 fl. drs. (3.70 to 11.09 c. c.)

**Tinctura Catechu Composita—Compound Tincture of Catechu.**DOSE:  $\frac{1}{2}$  to 3 fl. drs. (1.85 to 11.09 c. c.)**Tinctura Chiratæ—Tincture of Chirata.**DOSE:  $\frac{1}{2}$  to 2 fl. drs. (1.85 to 7.39 c. c.)**Tinctura Cimicifugæ—Tincture of Cimicifuga.**

DOSE: 1 to 2 fl. drs. (3.70 to 7.39 c. c.)

**Tinctura Cinchonæ—Tincture of Cinchona.**

DOSE: 1 to 2 fl. drs. (3.70 to 7.39 c. c.)

**Tinctura Cinchonæ Composita—Compound Tincture of Cinchona.**

DOSE: 1 to 4 fl. drs. (3.70 to 14.79 c. c.)

**Tinctura Cinnamomi—Tincture of Cinnamon.**

DOSE: 1 to 4 fl. drs. (3.70 to 14.79 c. c.)

**Tinctura Colchici Seminis—Tincture of Colchicum Seed—Tinctura Colchici, Pharm. 1880.**

DOSE: 10 to 60 min. (0.62 to 3.70 c. c.)

**Tinctura Croci—Tincture of Saffron.**

DOSE: 1 to 3 fl. drs. (3.70 to 11.09 c. c.)

**Tinctura Cubebæ—Tincture of Cubeb.**DOSE:  $\frac{1}{2}$  to 2 fl. drs. (1.85 to 7.39 c. c.)**Tinctura Digitalis—Tincture of Digitalis.**

DOSE: 5 to 30 min. (0.31 to 1.85 c. c.)

**Tincturæ Herbarum Recentium—Tinctures of Fresh Herbs.**

Fresh Herb, 50 gms. to 100 c. c.

**TRY MELLIN'S FOOD**  
**In Cholera Infantum.**

**Tinctura Ferri Chloridi**—*Tincture of Ferric Chloride.*

A hydro-alcoholic solution of Ferric Chloride, containing about 13.6% of the anhydrous salt, and corresponding to about 4.7% (4.69) of metallic iron.

Solution of Ferric Chloride, 25%.

DOSE: 5 to 20 min. (0.31 to 1.23 c. c.)

Prep: Liquor Ferri et Ammonii Acetatis.

**Tinctura Gallæ**—*Tincture of Nutgall.*

DOSE:  $\frac{1}{2}$  to 2 fl. drs. (1.85 to 7.39 c. c.)

**Tinctura Gelsemii**—*Tincture of Gelsemium.*

DOSE: 2 to 15 min. (0.12 to 0.92 c. c.)

**Tinctura Gentianæ Composita**—*Compound Tincture of Gentian.*

DOSE: 1 to 2 fl. drs. (3.70 to 7.39 c. c.)

**Tinctura Guaiaci**—*Tincture of Guaiac.*

DOSE:  $\frac{1}{2}$  to 1 fl. dr. (1.85 to 3.70 c. c.)

**Tinctura Guaiaci Ammoniata**—*Ammoniated Tincture of Guaiac.*

DOSE:  $\frac{1}{2}$  to 1 fl. dr. (1.85 to 3.70 c. c.)

**Tinctura Humuli**—*Tincture of Hops.*

DOSE: 1 to 8 fl. drs. (3.70 to 11.09 c. c.)

**Tinctura Hydrastis**—*Tincture of Hydrastis.*

DOSE:  $\frac{1}{2}$  to 1 fl. dr. (1.85 to 3.70 c. c.)

**Tinctura Hyoscyami**—*Tincture of Hyoscyamus.*

DOSE: 10 to 60 min. (0.62 to 3.70 c. c.)

**Tinctura Iodi**—*Tincture of Iodine.*

Iodine, 7 gms. in 100 c. c.

DOSE: 2 to 6 min. (0.12 to 0.37 c. c.)

**Tinctura Ipecacuanhæ et Opii**—*Tincture of Ipecac and Opium.*

DOSE: 5 to 15 min. (0.81 to 0.92 c. c.)

**Tinctura Kino**—*Tincture of Kino.*

DOSE:  $\frac{1}{2}$  to 2 fl. drs. (1.85 to 7.39 c. c.)

**Tinctura Krameriaæ**—*Tincture of Krameria.*

DOSE:  $\frac{1}{2}$  to 2 fl. drs. (1.85 to 7.39 c. c.)

**Tinctura Lactucarii**—*Tincture of Lactucarium.*

DOSE: 10 to 60 min. (0.62 to 3.70 c. c.)

Prep: Syrupus Lactucarii.

**Tinctura Lavandulæ Composita**—*Compound Tincture of Lavender*—Compound Spirit of Lavender.

DOSE:  $\frac{1}{2}$  to 2 fl. drs. (1.85 to 7.39 c. c.)

---

**TRY MELLIN'S FOOD**  
In Intestinal Indigestion.

**Tinctura Lobeliæ—Tincture of Lobelia.**

DOSE: 8 to 15 min.; 30 to 60 min. (0.49 to 0.92 c. c.; 1.85 to 3.70 c. c.)

**Tinctura Matico—Tincture of Matico.**

DOSE: 1 to 2 fl. drs. (3.70 to 7.39 c. c.)

**Tinctura Moschi—Tincture of Musk.**DOSE:  $\frac{1}{2}$  to 2 fl. drs. (1.85 to 7.39 c. c.)**Tinctura Myrrhæ—Tincture of Myrrh.**

DOSE: 15 to 60 min. (0.92 to 1.85 c. c.)

**Tinctura Nucis Vomicae—Tincture of Nux Vomica.**

0.3 gm. alkaloid in 100 c. c.

DOSE: 5 to 15 min. (0.31 to 0.92 c. c.)

**Tinctura Opii—Tincture of Opium—Laudanum.**

1.3 to 1.5 gm. Morphine in 100 c. c.)

DOSE: 5 to 15 min. (0.31 to 0.92 c. c.)

**Tinctura Opii Camphorata—Camphorated Tincture of Opium—Paregoric.**

DOSE: 1 to 4 fl. drs. (3.70 to 14.79 c. c.)

**Tinctura Opii Deodorati—Tincture of Deodorized Opium—Tinctura Opii Deodorata, Pharm. 1880.**

1.3 to 1.5 gm. Morphine in 100 c. c.

DOSE: 5 to 15 min. (0.31 to 0.92 c. c.)

Prep: Tinctura Ipecacuanhæ et Opii.

**Tinctura Physostigmatis—Tincture of Physostigma.**

DOSE: 5 to 10 min. (0.31 to 0.62 c. c.)

**Tinctura Pyrethri—Tincture of Pyrethrum.**  
External.**Tinctura Quassia—Tincture of Quassia.**DOSE:  $\frac{1}{2}$  to 1 fl. dr. (1.85 to 3.70 c. c.)**Tinctura Quillajæ—Tincture of Quillaja.**

DOSE: 15 to 30 min. (0.92 to 1.85 c. c.)

**Tinctura Rhei—Tincture of Rhubarb.**

DOSE: 1 to 4 fl. drs. (3.70 to 14.79 c. c.)

**Tinctura Rhei Aromatica—Aromatic Tincture of Rhubarb.**DOSE:  $\frac{1}{2}$  to 3 fl. drs. (1.85 to 11.09 c. c.)

Prep: Syrupus Rhei Aromaticus.

**Tinctura Rhei Dulcis—Sweet Tincture of Rhubarb.**

DOSE: 1 to 2 fl. drs. (3.70 to 7.39 c. c.)

**Tinctura Sanguinariæ—Tincture of Sanguinaria.**

DOSE: 15 to 30 min. (0.92 to 1.85 c. c.)

---

**TRY MELLIN'S FOOD**  
As a diet for Nursing Mothers.

**Tinctura Scillæ**—*Tincture of Squill.*

DOSE: 5 to 20 min. (0.31 to 1.23 c. c.)

**Tinctura Serpentariæ**—*Tincture of Serpentaria.*DOSE:  $\frac{1}{2}$  to 2 fl. drs. (1.85 to 7.39 c. c.)**Tinctura Stramonii Seminis**—*Tincture of Stramonium Seed* — *Tinctura Stramonii*, Pharm. 1880.

DOSE: 5 to 10 min. (0.31 to 0.62 c. c.)

**Tinctura Strophanthi** — *Tincture of Strophanthus.*

DOSE: 3 to 10 min. (0.18 to 0.62 c. c.)

**Tinctura Sumbul**—*Tincture of Sumbul.*

DOSE: 15 to 60 min. (0.92 to 3.70 c. c.)

**Tinctura Tolutana**—*Tincture of Tolu.*DOSE:  $\frac{1}{2}$  to 2 fl. drs, (1.85 to 7.39 c. c.)**Tinctura Valerianæ** *Tincture of Valerian.*

DOSE: 1 to 2 fl. drs. (3.70 to 7.39 c. c.)

**Tinctura Valerianæ Ammoniata**—*Ammoniated Tincture of Valerian.*DOSE:  $\frac{1}{2}$  to 1 fl. dr. (1.85 to 3.70 c. c.)**Tinctura Vanillæ**—*Tincture of Vanilla.*

Flavoring.

**Tinctura Veratri Viridis**—*Tincture of Veratrum Viride.*

DOSE: 1 to 5 min. (0.06 to 0.31 c. c.)

**Tinctura Zingiberis**—*Tincture of Ginger.*

DOSE: 15 to 60 min. (0.92 to 3.70 c. c.)

**Tragacantha**—*Tragacanth.*A gummy exudation from *Astragalus gummifer* and from other species of *Astragalus*.

Prep: Mucilago Tragacanthæ.

**Triticum**—*Triticum*—Couch grass.The rhizome of *Agropyrum repens* gathered in the spring and deprived of the roots.

DOSE: 1 to 8 drs. (3.888 to 31.103 gms.)

Prep: Extractum Tritici Fluidum.

**Triturationes**—*Triturations.*

Unless otherwise directed, Triturations are to be prepared by the following formula:

10% of the Substance.

90% Sugar of Milk.

**Trituratio Elaterini**—*Trituration of Elaterin.*

Elaterin, 10%.

DOSE:  $\frac{1}{2}$  to  $\frac{1}{4}$  gr. (0.0324 to 0.0405 gm.)**Trochisci Acidi Tannici**—*Troches of Tannic Acid.*

Tannic Acid, 0.06 gm. to each troche.

DOSE: 1 to 2 troches.

**MELLIN'S FOOD**

Makes cow's milk easily digestible.

**Trochisci Ammonii Chloridi—Troches of Ammonium Chloride.**

Ammonium Chloride, 0.10 gm. to each troche.

DOSE. 1 to 2 troches.

**Trochisci Catechu—Troches of Catechu.**

Catechu, 0.06 gm. to each troche.

DOSE: 1 to 2 troches.

**Trochisci Cretæ—Troches of Chalk.**

Prepared Chalk, 0.25 gm. to each troche.

DOSE. 1 to 5 troches.

**Trochisci Cubebæ—Troches of Cubeb.**

Oleoresin of Cubeb, 0.04 gm. to each troche.

DOSE: 1 to 2 troches.

**Trochisci Ferri—Troches of Iron.**

Ferric Hydrate, 0.30 gm. to each troche.

DOSE: 1 to 6 troches.

**Trochisci Glycyrrhizæ et Opii—Troches of Glycyrrhiza and Opium.**

Extract Glycyrrhiza, 0.15 gm. and Powdered Opium 0.005 to each troche.

DOSE: 1 to 2 troches.

**Troches Ipecacuanhæ—Troches of Ipecac.**

Ipecac, 0.02 gm. to each troche.

DOSE: 1 to 2 troches.

**Trochisci Kramerizæ—Troches of Krameria.**

Extract of Krameria, 0.06 gm. to each troche.

DOSE: 1 to 2 troches.

**Trochisci Menthæ Piperitæ—Troches of Peppermint.**

Oil of Peppermint, 0.01 c. c. to each troche.

DOSE. 1 to 5 troches.

**Trochisci Morphinæ et Ipecacuanhæ—Troches of Morphine and Ipecac.**

Morphine Sulphate, 0.0016 gm., and Ipecac, 0.005 gm. to each troche.

DOSE. 1 to 2 troches.

**Trochisci Potassii Chloratis—Troches of Potassium Chlorate.**

Potassium Chlorate, 0.30 gm. to each troche.

DOSE: 1 to 5 troches.

**Trochisci Santonini—Troches of Santonin.**

Santonin, 0.03 gm. to each troche.

DOSE: 1 to 6 troches.

**Trochisci Sodii Bicarbonatis—Troches of Sodium Bicarbonate.**

Sodium Bicarbonate, 0.20 gm. to each troche.

DOSE. 1 to 6 troches.

---

**MELLIN'S FOOD**  
Is readily assimilated.

**Trochisci Zingiberis—Troches of Ginger.**

Tincture of Ginger, 0.20 c. c. to each troche.

DOSE: 1 to 6 troches.

**Ulmus—Elm—Slippery Elm.**The inner bark of *Ulmus fulva*.

Prep. Mucilago Ulmi.

**Unguentum Ointment.**

Lard, 80%, Yellow Wax, 20%

**Unguentum Acidi Carbolici—Ointment of Carbolic Acid.**

Carbolic Acid, 5%.

**Unguentum Acidi Tannici—Ointment of Tannic Acid.**

Tannic Acid, 20%.

**Unguentum Aquæ Rosæ—Ointment of Rose Water—Cold Cream.****Unguentum Belladonnæ—Belladonna Ointment.**

Alcoholic Extract Belladonna Leaves, 10%

**Unguentum Chrysarobini—Chrysarobin Ointment.**

Chrysarobin, 5%.

**Unguentum Diachylon—Diachylon Ointment.**

L. ad Plaster, 50%.

**Unguentum Gallæ—Nutmeg Ointment.**

Nutmeg, 20%.

**Unguentum Hydrargyri—Mercurial Ointment—Blue Ointment.**

Mercury, 50%. Oleate Mercury, 2%.

**Unguentum Hydrargyri Ammoniatum—Ointment of Ammoniated Mercury.**

Ammoniated Mercury, 10%.

**Unguentum Hydrargyri Nitratis—Ointment of Mercuric Nitrate—Citrine Ointment.**

Mercury, about 7%.

**Unguentum Hydrargyri Oxidi Flavi—Ointment of Yellow Mercuric Oxide.**

Yellow Mercuric Oxide, 10%.

**Unguentum Hydrargyri Oxidi Rubri—Ointment of Red Mercuric Oxide.**

Red Mercuric Oxide, 10%.

**Unguentum Iodi—Iodine Ointment.**

Iodine, 4%. Potassium Iodide, 1%.

**Unguentum Iodoformi—Iodoform Ointment.**

Iodoform, 10%.

**MELLIN'S FOOD**

Is a soluble dry extract from wheat and malt.



**Unguentum Picis Liquidæ—Tar Ointment.**

Tar, 50%.

**Unguentum Plumbi Carbonatis—Ointment of Lead Carbonate.**

Lead Carbonate, 10%.

**Unguentum Plumbi Iodidi—Ointment of Lead Iodide.**

Lead Iodide, 10%.

**Unguentum Potassii Iodidi—Ointment of Potassium Iodide.**

Potassium Iodide, 12%.

**Unguentum Stramonii—Stramonium Ointment.**

Extract of Stramonium Seed, 10%.

**Unguentum Sulphuris—Sulphur Ointment.**

Washed Sulphur, 30%.

**Unguentum Veratrinæ—Veratrine Ointment.**

Veratrine, 4%.

**Unguentum Zinci Oxidi—Ointment of Zinc Oxide**

Zinc Oxide, 20%.

**Uva Ursi Uva Ursi—Bearberry.**The leaves of *Arctostaphylos Uva-ursi*.DOSE  $\frac{1}{4}$  to 1 dr (0.972 to 3.888 gms.)

Preps Extractum Uvæ Ursi. Extractum Uvæ Ursi Fluidum.

**Valeriana—Valerian.**The rhizome and roots of *Valeriana officinalis*.

DOSE: 10 to 30 grs. (0.648 to 1.944 gm.)

Preps. Extractum Valerianæ Fluidum. Tinctura Valerianæ. Tinctura Valerianæ Ammoniata.

**Vanilla—Vanilla.**The fruit of *Vanilla planifolia*.

Prep: Tinctura Vanillæ.

**Veratrina—Veratrine.**A mixture of alkaloids obtained from the seed of *Asagrea officinalis*.

DOSE. 1 30 to 1 10 gr. (0.0021 to 0.0064 gm.)

Preps Oleatum Veratrinæ. Unguentum Veratrinæ.

**Veratrum Viride—Veratrum Viride—American Hellebore.**The rhizome and roots of *Veratrum viride*.

DOSE. 1 to 5 grs. (0.0648 to 0.324 gm.)

Preps: Extractum Veratri Viridis Fluidum. Tinctura Veratri Viridis.

---

**MELLIN'S FOOD****Is entirely free from starch and cane sugar.**

**Viburnum Opulus — Viburnum Opulus—**  
**Cran p lark.**

The bark of *Viburnum Opulus*.

DOSE: 1 to 2 drs. (3.888 to 7.776 gms.)

Prep: *Extractum Viburni Opuli Fluidum*.

**Viburnum Prunifolium—Black Haw.**

The bark of *Virburnum prunifolium*.

DOSE.  $\frac{1}{4}$  to 1 dr. (0.972 to 3.888 gms.)

Prep. *Extractum Virburni Prunifolii Fluidum*.

**Vinum Album—White Wine.**

An alcoholic liquid made by fermenting the juice of fresh grapes, the fruit of *Vitis vinifera*, freed from seeds, stems, and skins.

10 to 14% by weight of absolute alcohol.

**Vinum Antimonii—Wine of Antimony.**

DOSE 10 to 30 min.,  $\frac{1}{2}$  to 2 fl. drs. (0.62 to 1.85 c. c.; 1.85 to 7.39 c. c.)

**Vinum Colchici Radicis—Wine of Colchicum Root.**

DOSE. 5 to 20 min. (0.31 to 1.23 c. c.)

**Vinum Colchici Seminis—Wine of Colchicum Seed.**

DOSE: 10 to 60 min. (0.62 to 3.70 c. c.)

**Vinum Ergotæ—Wine of Ergot.**

DOSE: 1 to 5 fl. drs. (3.70 to 11.00 c. c.)

**Vinum Ferri Amarum—Bitter Wine of Iron.**

DOSE: 1 to 3 fl. drs. (3.70 to 11.09 c. c.)

**Vinum Ferri Citratis—Wine of Ferric Citrate.**

DOSE: 1 to 2 fl. drs. (3.70 to 7.39 c. c.)

**Vinum Ipecacuanhæ—Wine of Ipecac.**

Fluid Extract Ipecac, 10%.

DOSE: 5 to 10 min. (0.31 to 0.62 c. c.)

**Vinum Opii—Wine of Opium.**

1.3 to 1.5 gm. Morphine in 100 c. c.

DOSE: 5 to 15 min. (0.31 to 0.92 c. c.)

**Vinum Rubrum—Red Wine.**

An alcoholic liquid made by fermenting the juice of fresh colored grapes, the fruit of *Vitis vinifera*, in presence of their skins.

10 to 14% by weight of absolute alcohol.

**Vitellus—Yolk of Egg.**

The yolk of the egg of *Gallus Bankiva*, var. *domestica*.

Prep: *Glyceritum Vitelli*.

**MELLIN'S FOOD**

Stimulates the bowels to regular action.

**Xanthoxylum—Xanthoxylum—Prickly Ash.**

The bark of *Xanthoxylum americanum* and of *Xanthoxylum Clava-Herculis*.

DOSE: 10 to 30 grs. (0.648 to 1.944 gm.)

Prep: Extractum Xanthoxyli Fluidum.

**Zea—Zea—Corn-Silk.**

The styles and stigmas of *Zea Mays*.

Prep: Extractum Zeæ Fluidum.

**Zinci Acetas—Zinc Acetate.**

DOSE:  $\frac{1}{2}$  to 2 grs. (0.0324 to 0.1296 gm.)

**Zinci Bromidum—Zinc Bromide.**

DOSE: 1 to 2 grs. (0.0648 to 0.1296 gm.)

**Zinci Carbonas Præcipitatus—Precipitated Zinc Carbonate.**

Externally.

**Zinci Chloridum—Zinc Chloride.**

Externally.

**Zinci Iodidum—Zinc Iodide.**

DOSE: 1 to 2 grs. (0.0648 to 0.1296 gms.)

**Zinci Oxidum—Zinc Oxide.**

DOSE: 1 to 5 grs. (0.0648 to 0.324 gm.)

Prep: Unguentum Zinci Oxidi.

**Zinci Phosphidum—Zinc Phosphide.**

DOSE: 1-16 to  $\frac{1}{4}$  gr. (0.0040 to 0.0216 gm.)

**Zinci Sulphas—Zinc Sulphate.**

DOSE: 1 to 3 grs.; 10 to 60 grs. (0.0648 to 0.1944 gm.; 0.648 to 3.888 gms.)

**Zinci Valerianas—Zinc Valerianate.**

DOSE:  $\frac{1}{2}$  to 2 grs. (0.0324 to 0.1296 gm.)

**Zincum—Zinc.**

Metallic zinc in the form of thin sheets, or in irregular, granulated pieces, or moulded into thin pencils, or in a state of fine powder.

**Zingiber—Ginger.**

The rhizome of *Zingiber officinale*.

DOSE: 5 to 15 grs. (0.324 to 0.972 gm.)

Preps: Extractum Zingiberis Fluidum. Oleoresina Zingiberis. Pulvis Aromaticus. Pulvis Rhei Compositus. Tinctura Zingiberis.

---

"With children sick with Typhoid Fever, I have found MELLIN'S an excellent food."

# MELLIN'S FOOD

—FOR—

INFANTS AND INVALIDS.

---

**Mellin's Food** when dissolved in water and added to fresh cow's milk, produces Liebig's Food for infants, also called Liebig's Soup.

**Mellin's Food** added to fresh cow's milk supplies carbohydrates, albuminates and salts to make up the deficiency of these constituents in cow's milk as compared with mother's milk.

**Mellin's Food** added to fresh cow's milk furnishes alkali sufficient not only for permanent reaction, but also for the conversion of the caseine of the milk into soluble albuminates, thus changing the character of the cow's milk in such a manner as to assure its easy digestion by infants, and by invalids and by delicate persons and by persons of weak digestion.

---

DOLIBER-GOODALE CO.,

BOSTON, MASS.







